Background characteristics of respondents

Percent distribution of background characteristics of women age 15 to 49

Background characteristics	Urban	Rural	Percentage of sample	Weighted number	Unweighted number
Age group					
15–19	46.0	54.0	100.0	23.3	392
20–24	40.3	59.7	100.0	18.8	317
25–29	37.3	62.7	100.0	16.4	277
30–34	37.5	62.5	100.0	14.7	248
35–39	47.2	52.8	100.0	11.5	194
40–44	45.0	55.0	100.0	9.7	164
45–49	47.1	52.9	100.0	5.5	92
Marital status					
Married	33.6	66.4	100.0	77.0	1,297
Living together	(71.0)	(29.0)	(100.0)	0.2	3
Divorced	53.6	46.4	100.0	2.3	39
Widowed	75.0	25.0	100.0	1.5	25
Never married	73.9	26.1	100.0	18.9	319
Parity					
None	58.9	41.1	100.0	27.8	469
1-2	31.8	68.2	100.0	20.1	338
3-4	35.1	64.9	100.0	17.5	295
5 or more	38.8	61.2	100.0	34.6	582
Education					
Never attended	15.7	84.3	100.0	45.4	765
Primary	32.5	67.5	100.0	20.8	350
Secondary	82.2	17.8	100.0	25.0	420
Higher	89.9	10.1	100.0	8.8	148
Wealth quintile					
Lowest	0.6	99.4	100.0	18.0	302
Lower	3.0	97.0	100.0	17.6	297
Middle	27.0	73.0	100.0	18.0	303
Higher	71.3	28.7	100.0	22.9	385
Highest	87.4	12.6	100.0	23.5	396
Total	**42.4**	**57.6**	**100.0**	**100.0**	**1,684**

Current use of contraception by background characteristics

Percentage of women ages 15 to 49 currently using a contraceptive method, by type and background characteristics

Background characteristic	Any method		Modern method[1]		Traditional method	
	All women	Married[2] women	All women	Married[2] women	All women	Married[2] women
Number	1,684	1,300	1,684	1,300	1,684	1,300
Total	**5.6**	**7.1**	**4.9**	**6.1**	**0.8**	**1.0**
Age group						
15–19	1.2	3.1	1.2	3.1	0.0	0.0
20–24	3.0	3.9	2.1	2.7	0.9	1.2
25–29	7.9	7.4	6.7	6.2	1.2	1.3
30–34	8.1	8.5	6.9	7.2	1.2	1.3
35–39	8.1	8.8	7.9	8.5	0.2	0.3
40–44	10.3	11.0	9.6	10.3	0.6	0.7
45–49	6.5	7.2	3.9	4.3	2.6	2.9
Marital status						
Married[2]	7.1	na	6.1	na	1.0	na
Unmarried, sexually active	(0.0)	na	(0.0)	na	(0.0)	na
Parity						
0–1	1.5	2.2	1.5	2.2	0.0	0.0
2–3	7.9	8.3	6.5	6.9	1.4	1.4
4 or more	8.1	8.5	6.9	7.3	1.1	1.2
Residence						
Urban	10.3	16.2	9.0	14.0	1.4	2.2
Rural	2.2	2.4	1.8	2.1	0.3	0.4
Education						
Never attended	2.5	2.8	2.3	2.5	0.3	0.3
Primary	6.2	7.0	4.8	5.5	1.4	1.6
Secondary	6.6	13.3	6.0	12.0	0.6	1.3
Higher	17.0	27.5	14.5	23.2	2.5	4.3
Wealth quintile						
Lowest	0.3	0.4	0.0	0.0	0.3	0.4
Lower	0.7	0.8	0.3	0.4	0.3	0.4
Middle	4.4	5.3	3.7	4.4	0.8	0.9
Higher	7.7	10.9	7.0	9.8	0.7	1.1
Highest	12.3	19.8	10.8	17.4	1.5	2.4

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

Contraceptive method mix by background characteristics

Percent distribution of current contraceptive users ages 15 to 49, by method type, marital status and background characteristics

Background characteristics	Sterilization	Implants	IUD	Inject-ables	Pill	Emergency contraception	Condom	Other modern	Other traditional	Total
All users Number = 95										
Total	**0.0**	**14.0**	**6.3**	**39.9**	**20.0**	**3.0**	**12.0**	**3.3**	**11.6**	**100.0**
Age group										
15–19	*(0.0)*	*(0.0)*	*(0.0)*	*(52.4)*	*(47.6)*	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	*(100.0)*
20–24	(0.0)	(17.8)	(0.0)	(35.1)	(16.9)	(0.0)	(0.0)	(0.0)	(30.2)	(100.0)
25–29	(0.0)	(26.7)	(0.0)	(32.8)	(6.7)	(12.9)	(5.9)	(0.0)	(15.0)	(100.0)
30–34	(0.0)	(11.8)	(13.7)	(39.8)	(16.5)	(0.0)	(3.1)	(5.1)	(10.0)	(100.0)
35–39	(0.0)	(4.1)	(9.5)	(55.0)	(28.4)	(0.0)	(0.0)	(0.0)	(3.0)	(100.0)
40–44	(0.0)	(16.5)	(6.1)	(43.2)	(28.0)	(0.0)	(0.0)	(6.1)	(0.0)	(100.0)
45–49	(0.0)	(0.0)	(10.8)	(14.0)	(17.3)	(0.0)	(0.0)	(17.8)	(40.1)	(100.0)
Marital status										
Married[1]	0.0	14.4	6.5	41.1	20.6	0.0	2.1	3.4	11.9	100.0
Unmarried, sexually active	na	na	na	na	na	na	na	na	na	na
Parity										
0–1	(0.0)	(26.0)	(0.0)	(28.3)	(14.4)	(31.3)	(0.0)	(0.0)	(0.0)	(100.0)
2–3	0.0	23.8	6.1	29.1	16.8	0.0	6.8	0.0	17.3	100.0
4 or more	0.0	7.4	7.4	46.9	22.4	0.0	0.0	5.4	10.6	100.0
Residence										
Urban	0.0	15.1	8.1	35.4	22.1	3.8	0.9	4.2	10.4	100.0
Rural	(0.0)	(10.2)	(0.0)	(55.6)	(12.3)	(0.0)	(6.1)	(0.0)	(15.8)	(100.0)
Education										
Never attended	(0.0)	(0.0)	(0.0)	(61.3)	(28.3)	(0.0)	(0.0)	(0.0)	(10.4)	(100.0)
Primary	(0.0)	(0.0)	(7.8)	(50.8)	(19.3)	(0.0)	(0.0)	(4.8)	(17.4)	(100.0)
Secondary	0.0	15.7	0.0	45.0	24.6	5.7	0.0	0.0	9.0	100.0
Higher	0.0	35.4	17.0	9.2	9.3	4.8	5.1	8.3	10.7	100.0
Wealth quintile										
Lowest	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	*(100.0)*	*(100.0)*
Lower	*(0.0)*	*(0.0)*	*(0.0)*	*(49.5)*	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	*(50.5)*	*(100.0)*
Middle	*(0.0)*	*(0.0)*	*(0.0)*	*(65.8)*	*(17.1)*	*(0.0)*	*(0.0)*	*(0.0)*	*(17.1)*	*(100.0)*
Higher	0.0	0.0	2.2	49.3	31.5	5.3	2.1	0.0	9.6	100.0
Highest	0.0	27.3	10.9	27.5	14.9	2.5	2.6	6.4	7.8	100.0

[1]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

| Background characteristics	Sterilization	Implants	IUD	Inject-ables	Pill	Emergency contraception	Condom	Other modern	Other traditional	Total
Married users										
Number = 92										
Total	**0.0**	**14.4**	**6.5**	**41.1**	**20.6**	**0.0**	**2.1**	**3.4**	**11.9**	**100.0**
Age group										
15–19	(0.0)	(0.0)	(0.0)	(52.4)	(47.6)	(0.0)	(0.0)	(0.0)	(0.0)	(100.0)
20–24	(0.0)	(17.8)	(0.0)	(35.1)	(16.9)	(0.0)	(0.0)	(0.0)	(30.2)	(100.0)
25–29	(0.0)	(30.6)	(0.0)	(37.7)	(7.7)	(0.0)	(6.8)	(0.0)	(17.2)	(100.0)
30–34	(0.0)	(11.8)	(13.7)	(39.8)	(16.5)	(0.0)	(3.1)	(5.1)	(10.0)	(100.0)
35–39	(0.0)	(4.1)	(9.5)	(55.0)	(28.4)	(0.0)	(0.0)	(0.0)	(3.0)	(100.0)
40–44	(0.0)	(16.5)	(6.1)	(43.2)	(28.0)	(0.0)	(0.0)	(6.1)	(0.0)	(100.0)
45–49	(0.0)	(0.0)	10.8	(14.0)	(17.3)	(0.0)	(0.0)	(17.8)	(40.1)	(100.0)
Marital status										
Married[1]	na	na	na	na	na	na	na	na	na	na
Unmarried, sexually active	na	na	na	na	na	na	na	na	na	na
Parity										
0–1	(0.0)	(37.8)	(0.0)	(41.2)	(21.0)	(0.0)	(0.0)	(0.0)	(0.0)	(100.0)
2–3	0.0	23.8	6.1	29.1	16.8	0.0	6.8	0.0	17.3	100.0
4 or more	0.0	7.4	7.4	46.9	22.4	0.0	0.0	5.4	10.6	100.0
Residence										
Urban	0.0	15.7	8.4	36.8	23.0	0.0	0.9	4.4	10.8	100.0
Rural	(0.0)	(10.2)	(0.0)	(55.6)	(12.3)	(0.0)	(6.1)	(0.0)	(15.8)	(100.0)
Education										
Never attended	(0.0)	(0.0)	(0.0)	(61.3)	(28.3)	(0.0)	(0.0)	(0.0)	(10.4)	(100.0)
Primary	(0.0)	(0.0)	(7.8)	(50.8)	(19.3)	(0.0)	(0.0)	(4.8)	(17.4)	(100.0)
Secondary	0.0	16.7	0.0	47.7	26.1	0.0	0.0	0.0	9.5	100.0
Higher	0.0	37.2	17.9	9.7	9.8	0.0	5.4	8.8	11.3	100.0
Wealth quintile										
Lowest	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(100.0)	(100.0)
Lower	(0.0)	(0.0)	(0.0)	(49.5)	(0.0)	(0.0)	(0.0)	(0.0)	(50.5)	(100.0)
Middle	(0.0)	(0.0)	(0.0)	(65.8)	(17.1)	(0.0)	(0.0)	(0.0)	(17.1)	(100.0)
Higher	0.0	0.0	2.3	52.0	33.2	0.0	2.3	0.0	10.2	100.0
Highest	0.0	28.0	11.2	28.2	15.3	0.0	2.7	6.6	8.0	100.0

[1]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

PMA2020 Performance Monitoring and Accountability 2020

© 2017 Performance Monitoring and Accountability 2020 (PMA2020)

Current Method Mix Among **All and Married** Contraceptive Users (Age 15-49)

Number of users:
95

Number of users:
92

Percentage

	All	Married
Other traditional	11.6	11.9
Other modern	3.3	3.4
Male condoms	2.0	2.1
Emergency contraception	3.0	
Pills	20.0	20.6
Injectables	39.9	41.1
Implants	14.0	14.4
IUD	6.3	6.5

Legend:
- Other traditional
- Other modern
- Emergency contraception
- Male condoms
- Pills
- Injectables
- Implants
- IUD
- Female Sterilization

PMA2020 Performance Monitoring and Accountability 2020

© 2017 Performance Monitoring and Accountability 2020 (PMA2020)

Unmet need for family planning

Percentage of women ages 15 to 49 with unmet need for family planning (for spacing or limiting), by marital status and background characteristics

Background characteristics	All women			Married women		
Number	**1,684**			**1,300**		
	For spacing	**For limiting**	**Total**	**For spacing**	**For limiting**	**Total**
Total	**25.5**	**4.7**	**30.2**	**32.2**	**6.0**	**38.2**
Age group						
15–19	14.8	0.0	14.8	33.9	0.0	33.9
20–24	25.7	1.0	26.8	32.5	1.4	33.9
25–29	39.9	1.3	41.2	43.3	0.8	44.1
30–34	34.2	5.5	39.7	35.9	5.8	41.6
35–39	29.4	11.9	41.3	31.0	12.8	43.7
40–44	19.6	13.7	33.3	21.0	14.7	35.8
45–49	6.2	14.1	20.3	6.9	15.7	22.6
Marital status						
Married[1]	32.2	6.0	38.2	na	na	na
Unmarried, sexually active	(49.9)	(2.3)	(52.3)	na	na	na
Parity						
0–1	13.5	0.0	13.5	26.8	0.0	26.8
2–3	38.7	1.4	40.2	40.0	1.5	41.5
4 or more	29.3	10.3	39.6	30.6	10.6	41.2
Residence						
Urban	16.9	7.4	24.2	25.9	11.8	37.6
Rural	31.9	2.7	34.6	35.4	3.0	38.4
Education						
Never attended	32.7	4.4	37.1	34.8	4.7	39.5
Primary	27.4	7.6	35.1	31.0	8.4	39.3
Secondary	15.1	3.0	18.1	29.1	6.3	35.4
Higher	13.4	4.2	17.6	22.9	7.1	30.0
Wealth quintile						
Lowest	40.6	1.8	42.4	42.7	1.8	44.5
Lower	30.7	2.6	33.2	34.2	2.9	37.1
Middle	28.2	5.3	33.5	31.1	6.4	37.5
Higher	16.9	6.4	23.3	24.0	9.2	33.1
Highest	16.6	6.3	22.9	27.5	10.4	37.9

[1]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

PMA2020 · Performance Monitoring and Accountability 2020

© 2017 Performance Monitoring and Accountability 2020 (PMA2020)

Need and demand for family planning

Percentage women ages 15 to 49 with met need, unmet need, and total demand for family planning, and percentage of demand for family planning satisfied with modern methods, by marital status and background characteristics

Background characteristics	All women					Married women				
Number	**1,684**					**1,300**				
	Modern method[1]	Any method	Unmet need	Total demand	Demand satisfied by modern method	Modern method[2]	Any method	Unmet need	Total demand	Demand satisfied by modern method
Total	**4.9**	**5.6**	**30.2**	**35.8**	**13.5**	**6.1**	**7.1**	**38.2**	**45.2**	**13.4**
Age group										
15–19	1.2	1.2	14.8	16.1	7.6	3.1	3.1	33.9	37.0	8.5
20–24	2.1	3.0	26.8	29.7	6.9	2.7	3.9	33.9	37.8	7.2
25–29	6.7	7.9	41.2	49.0	13.6	6.2	7.4	44.1	51.6	11.9
30–34	6.9	8.1	39.7	47.9	14.4	7.2	8.5	41.6	50.2	14.4
35–39	7.9	8.1	41.3	49.5	16.0	8.5	8.8	43.7	52.5	16.2
40–44	9.6	10.3	33.3	43.5	22.1	10.3	11.0	35.8	46.8	22.1
45–49	3.9	6.5	20.3	26.8	14.5	4.3	7.2	22.6	29.8	14.5
Marital status										
Married[2]	6.1	7.1	38.2	45.2	13.4	na	na	na	na	na
Unmarried, sexually active	(0.0)	(0.0)	(52.3)	(52.3)	(0.0)	na	na	na	na	na
Parity										
0–1	1.5	1.5	13.5	15.0	9.7	2.2	2.2	26.8	29.0	7.6
2–3	6.5	7.9	40.2	48.1	13.6	6.9	8.3	41.5	49.8	13.8
4 or more	6.9	8.1	39.6	47.7	14.5	7.3	8.5	41.2	49.7	14.6
Residence										
Urban	9.0	10.3	24.2	34.6	26.0	14.0	16.2	37.6	53.8	26.0
Rural	1.8	2.2	34.6	36.8	4.9	2.1	2.4	38.4	40.9	5.0
Education										
Never attended	2.3	2.5	37.1	39.7	5.8	2.5	2.8	39.5	42.2	5.9
Primary	4.8	6.2	35.1	41.2	11.6	5.5	7.0	39.3	46.3	11.8
Secondary	6.0	6.6	18.1	24.7	24.4	12.0	13.3	35.4	48.7	24.7
Higher	14.5	17.0	17.6	34.6	41.8	23.2	27.5	30.0	57.5	40.4
Wealth quintile										
Lowest	0.0	0.3	42.4	42.8	0.0	0.0	0.4	44.5	44.8	0.0
Lower	0.3	0.7	33.2	33.9	1.0	0.4	0.8	37.1	37.8	1.0
Middle	3.7	4.4	33.5	37.9	9.7	4.4	5.3	37.5	42.8	10.3
Higher	7.0	7.7	23.3	31.0	22.5	9.8	10.9	33.1	44.0	22.2
Highest	10.8	12.3	22.9	35.2	30.7	17.4	19.8	37.9	57.7	30.1

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

Pregnancy intentions

Unintended births among women ages 15 to 49 with one or more births in the previous 5 years or who are currently pregnant: Percent distribution reporting last or current pregnancy was wanted then, wanted later or not wanted at all, by marital status and background characteristics

Background characteristics	All women with 1 or more births				Married women with 1 or more births			
Number	**1,005**				**974**			
	Wanted then	Wanted later	Not at all	Total	Wanted then	Wanted later	Not at all	Total
Total	**54.8**	**39.4**	**5.8**	**100.0**	**54.6**	**39.7**	**5.7**	**100.0**
Age group								
15–19	61.9	38.1	0.0	100.0	61.9	38.1	0.0	100.0
20–24	65.3	33.2	1.6	100.0	64.7	33.6	1.7	100.0
25–29	52.2	45.5	2.3	100.0	52.3	45.8	1.9	100.0
30–34	52.5	41.8	5.7	100.0	51.6	42.6	5.9	100.0
35–39	51.4	40.7	7.9	100.0	51.7	40.8	7.5	100.0
40–44	46.2	36.5	17.3	100.0	46.8	35.7	17.5	100.0
45–49	44.0	24.1	31.9	100.0	41.3	25.3	33.4	100.0
Marital status								
Married[1]	54.6	39.7	5.7	100.0	na	na	na	na
Unmarried, sexually active	(86.2)	(13.8)	(0.0)	(100.0)	na	na	na	na
Parity								
0–1	63.4	36.6	0.0	100.0	63.5	36.5	0.0	100.0
2–3	53.6	45.0	1.4	100.0	53.1	45.5	1.4	100.0
4 or more	53.0	37.1	9.9	100.0	52.8	37.4	9.8	100.0
Residence								
Urban	52.0	36.5	11.5	100.0	51.9	36.7	11.4	100.0
Rural	56.3	40.9	2.8	100.0	55.9	41.2	2.9	100.0
Education								
Never attended	59.4	35.5	5.1	100.0	59.2	35.6	5.2	100.0
Primary	47.5	44.4	8.0	100.0	46.8	45.3	7.9	100.0
Secondary	53.3	43.1	3.7	100.0	53.7	43.2	3.1	100.0
Higher	45.6	45.7	8.7	100.0	44.6	46.0	9.5	100.0
Wealth quintile								
Lowest	51.2	47.5	1.2	100.0	50.6	48.1	1.3	100.0
Lower	60.3	34.8	4.9	100.0	60.3	34.8	5.0	100.0
Middle	54.5	38.9	6.6	100.0	53.6	39.6	6.8	100.0
Higher	57.9	33.7	8.4	100.0	58.0	33.8	8.2	100.0
Highest	50.0	41.7	8.3	100.0	50.1	41.8	8.1	100.0

[1]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

PMA2020 | Performance Monitoring and Accountability 2020

Contraceptive choice

Percent distribution of women ages 15 to 49 who are currently using modern contraception[1] reporting on who decided on the contraceptive method, by background characteristics

Background characteristics	Self	Jointly with partner or provider	Provider alone/ partner alone/other	Total
Number	**28**	**43**	**23**	**95**
Total	**29.9**	**45.8**	**24.3**	**100.0**
Age group				
15–19	*(0.0)*	*(26.7)*	*(73.3)*	*(100.0)*
20–24	(46.1)	(40.3)	(13.7)	(100.0)
25–29	(39.6)	(50.2)	(10.2)	(100.0)
30–34	(5.0)	(52.7)	(42.3)	(100.0)
35–39	(39.5)	(52.1)	(8.5)	(100.0)
40–44	(42.1)	(26.4)	(31.4)	(100.0)
45–49	(17.8)	(68.1)	(14.0)	(100.0)
Marital status				
Married[2]	29.5	47.2	23.3	100.0
Unmarried, sexually active	na	na	na	na
Parity				
0–1	(13.6)	(40.4)	(46.0)	100.0
2–3	26.6	58.3	15.1	100.0
4 or more	34.0	40.5	25.5	100.0
Residence				
Urban	32.2	42.9	24.9	100.0
Rural	(21.9)	(55.9)	(22.2)	100.0
Education				
Never attended	(32.8)	(28.3)	(38.8)	(100.0)
Primary	(28.2)	(45.9)	(25.9)	(100.0)
Secondary	37.1	38.2	24.7	100.0
Higher	21.8	66.3	11.8	100.0
Wealth quintile				
Lowest	*(100.0)*	*(0.0)*	*(0.0)*	*(100.0)*
Lower	*(0.0)*	*(100)*	*(0.0)*	*(100.0)*
Middle	*(32.8)*	*(19.2)*	*(48.0)*	*(100.0)*
Higher	34.9	39.0	26.1	100.0
Highest	25.8	56.0	18.2	100.0

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Payment for family planning

Percentage of women ages 15 to 49 currently using a modern contraceptive method[1] who paid fees for services, by marital status and background characteristics

Background characteristics	All women who used in past 12 months	All married women who used in past 12 month
Number	**95**	**92**
Total	**50.6**	**52.2**
Age group		
15–19	*(55.9)*	*(55.9)*
20–24	(64.9)	(64.9)
25–29	(36.0)	(41.3)
30–34	(50.9)	(50.9)
35–39	(53.2)	(53.2)
40–44	(65.3)	(65.3)
45–49	(28.1)	(28.1)
Marital status		
Married[2]	52.2	na
Unmarried, sexually active	na	na
Parity		
0–1	(37.6)	(54.8)
2–3	37.2	37.2
4 or more	59.2	59.2
Residence		
Urban	52.1	54.1
Rural	(45.6)	(45.6)
Education		
Never attended	(69.9)	(69.9)
Primary	(43.1)	(43.1)
Secondary	62.3	66.1
Higher	30.4	31.9
Wealth quintile		
Lowest	*(0.0)*	*(0.0)*
Lower	*(50.5)*	*(50.5)*
Middle	*(55.8)*	*(55.8)*
Higher	68.3	72.2
Highest	39.5	40.5

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Method Information Index Components

Percentage of current modern[1] contraceptive users who reported whether provider informed them about other methods, side effects and, if informed of side effects, what to do, by marital status and background characteristics[2]

Background characteristics	Informed about other methods		Informed about side effects		Told what to do if experienced side effects	
	All users	Married users	All users	Married users	All users	Married users
Number	95	92	86	83	49	49
Total	**53.6**	**55.3**	**57.0**	**58.9**	**81.1**	**81.1**
Age group						
15–19	*(20.9)*	*(20.9)*	*(20.9)*	*(20.9)*	*(100)*	*(100)*
20–24	(36.0)	(36.0)	(67.3)	(67.3)	(74.1)	(74.1)
25–29	(60.7)	(69.6)	(61.3)	(71.6)	(75.4)	(75.4)
30–34	(59.1)	(59.1)	(53.7)	(53.7)	(88.4)	(88.4)
35–39	(61.7)	(61.7)	(71.4)	(71.4)	(77.3)	(77.3)
40–44	(51.9)	(51.9)	(43.2)	(43.3)	(96.1)	(96.1)
45–49	*(46.6)*	*(46.6)*	*(76.6)*	*(76.6)*	*(62.4)*	*(62.4)*
Marital status						
Married[3]	55.3	na	58.9	na	81.1	na
Unmarried, sexually active	na	na	na	na	na	na
Parity						
0–1	(14.4)	(21.0)	(26.0)	(37.8)	*(100)*	*(100)*
2–3	68.2	68.2	(77.0)	(78.0)	(75.6)	(75.6)
4 or more	52.6	52.6	52.1	52.1	83.6	83.6
Residence						
Urban	59.6	61.9	56.2	58.7	84.8	84.8
Rural	(32.8)	(32.8)	(59.5)	(59.5)	(69.6)	(69.6)
Education						
Never attended	(38.9)	(38.9)	(30.9)	(30.9)	(45.3)	(45.3)
Primary	(42.2)	(42.2)	(34.0)	(34.0)	(100)	(100)
Secondary	51.0	54.0	65.6	69.8	(84.6)	(84.6)
Higher	76.7	80.6	(91.0)	(96.5)	(83.4)	(83.4)
Wealth quintile						
Lowest	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	na	na
Lower	*(0.0)*	*(0.0)*	*(49.5)*	*(49.5)*	*(0.0)*	*(0.0)*
Middle	(20.4)	(20.4)	(29.4)	(29.4)	*(100)*	*(100)*
Higher	45.5	48.0	31.9	(33.9)	(59.3)	(59.3)
Highest	71.1	72.9	82.7	85.1	87.0	87.0

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2] Informed about other methods" asked of all current contraceptive users,"informed about side effects" asked of current modern contraceptive users, and "told what to do if experienced side effects" asked of current modern contraceptive users who indicated they were told about side effects.

[3]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Quality of care for family planning

Percentage of current modern[1] contraceptive users who would return, would refer a friend/relative, or would return and refer friend/relative to provider, by background characteristics

Background characteristics	Percent who would return to provider	Percent who would refer relative/friend	Percent who would return and refer friend/relative
Number	**86**	**86**	**86**
Total	**77.4**	**86.8**	**77.1**
Age group			
15–19	*(100.0)*	*(100.0)*	*(100.0)*
20–24	(67.3)	(78.5)	(67.3)
25–29	(77.2)	(80.5)	(77.2)
30–34	(78.5)	(88.1)	(78.5)
35–39	(76.1)	(95.4)	(76.1)
40–44	(70.4)	(81.5)	(68.7)
45–49	*(100.0)*	*(100.0)*	*(100.0)*
Marital status			
Married[2]	80.1	89.7	79.7
Unmarried, sexually active	na	na	na
Parity			
0–1	(68.7)	(68.7)	(68.7)
2–3	(86.4)	(89.6)	(86.4)
4 or more	74.6	88.5	74.1
Residence			
Urban	76.8	84.4	76.4
Rural	(79.5)	(94.8)	(79.5)
Education			
Never attended	(76.5)	(89.6)	(76.5)
Primary	(70.9)	(88.6)	(70.9)
Secondary	76.5	81.1	75.5
Higher	(84.2)	(89.4)	(84.2)
Wealth quintile			
Lowest	*(0.0)*	*(0.0)*	*(0.0)*
Lower	*(49.5)*	*(100.0)*	*(49.5)*
Middle	(86.3)	(93.2)	(86.3)
Higher	70.9	80.5	70.9
Highest	82.3	90.5	81.7

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Knowledge of family planning

Percentage of women ages 15 to 49 who reported receiving family planning information from a health worker in the past 12 months, by marital status and background characteristics

Background characteristics	Visited by health worker who discussed family planning		Visited facility and provider discussed family planning		Either visited by health worker or visited facility and discussed family planning	
	All women	Married women[1]	All women	Married women[1]	All women	Married women[1]
Number	**1,684**	**1,300**	**1,684**	**1,300**	**1,684**	**1,300**
Total	**7.4**	**8.5**	**13.2**	**16.0**	**17.3**	**20.4**
Age group						
15–19	3.2	6.0	2.5	5.5	5.1	10.0
20–24	7.3	8.1	15.9	18.4	20.2	22.7
25–29	10.6	10.4	21.4	23.0	26.6	27.4
30–34	8.0	8.4	15.9	16.7	20.8	21.8
35–39	12.1	11.6	18.8	19.5	23.5	24.0
40–44	8.0	7.8	9.9	9.2	14.2	13.8
45–49	4.2	3.2	11.9	10.2	13.8	12.4
Marital status						
Married[1]	8.5	na	16.0	na	20.4	na
Unmarried, sexually active	(0.0)	na	(14.6)	na	(14.6)	na
Parity						
0–1	4.1	6.4	6.5	12.3	9.3	15.9
2–3	11.0	10.8	16.2	16.1	22.1	21.5
4 or more	8.6	8.2	17.5	17.5	21.8	21.7
Residence						
Urban	11.4	15.7	16.5	24.7	21.8	31.5
Rural	4.5	4.8	10.8	11.6	14.0	14.8
Education						
Never attended	5.5	5.8	9.5	9.7	12.9	13.4
Primary	8.6	8.4	18.3	20.2	22.7	24.2
Secondary	10.7	19.3	13.5	26.9	19.6	36.2
Higher	5.0	5.7	18.8	27.2	20.1	28.0
Wealth quintile						
Lowest	2.7	2.8	5.3	4.8	7.0	6.7
Lower	3.1	3.4	11.5	12.9	13.7	15.4
Middle	6.1	6.9	13.4	15.3	17.9	20.3
Higher	12.2	15.3	13.7	19.1	20.6	27.5
Highest	10.7	15.3	20.0	30.2	24.1	35.0

[1]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

List of indicators for sampling errors, PMA2016/Kano

Variable	Estimation	Base population
Currently using a modern method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using a traditional method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using any contraceptive method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using injectables	Proportion	All women ages 15–49 Married women ages 15–49
Currently using male condoms	Proportion	All women ages 15–49 Married women ages 15–49
Currently using implants	Proportion	All women ages 15–49 Married women ages 15–49
Chose method by self or jointly in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49
Paid fees for family planning services in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49
Informed by provider about other methods	Proportion	All women ages 15–49 Married women ages 15–49
Informed by provider about side effects	Proportion	All women ages 15–49 Married women ages 15–49
Satisfied with provider: Would return and refer friend/relative to provider	Proportion	All women ages 15–49 Married women ages 15–49
Visited by health worker who talked about family planning information in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49

Characteristics of service delivery points

Number and percentage of service delivery points (SDPs) by background characteristics

Background characteristics	Number	Percentage
Total	44	100.0
SDP type		
Hospital	5	11.4
Health center[1]	20	45.5
Pharmacy[2]	18	40.9
Other	1	2.3
Sector		
Public	24	54.5
Private	20	45.5
Offers family planning		
No	10	22.7
Yes	34	77.3

[1]Includes Health Post, Health Clinic, and Maternity Center
[2]Includes Chemist and Patent Medicine Vendors

Charging fees for family planning

Percentage of service delivery points (SDPs) that charge fees for family planning services, by sector and background characteristics, among SDPs offering family planning

Background characteristic	Sector	
	Public	**Private**
Number of SDPs	17	17
Total	(11.8)	(88.2)
SDP type		
Hospital	*(0.0)*	na
Health center[1]	(15.4)	*(100.0)*
Pharmacy[2]	na	(87.5)
Other	na	na

[1]Includes Health Post, Health Clinic, and Maternity Center

[2]Includes Chemist and Patent Medicine Vendors

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Contraceptive choice: Availability of at least 3 or 5 modern contraceptive methods

Percentage of public and private service delivery points (SDPs) offering at least 3 or 5 modern contraceptive methods[1], by sector and background characteristics, among all SDPs

Background characteristic	3 or more methods		5 or more methods	
	Public	Private	Public	Private
Number of SDPs	24	20	24	20
Total	(54.2)	(15.0)	(29.2)	(5.0)
SDP type				
Hospital	(80.0)	na	(80.0)	na
Health Center[2]	(47.4)	*(100.0)*	(15.8)	*(0.0)*
Pharmacy[3]	na	(11.1)	na	(5.6)
Other	na	*(0.0)*	na	*(0.0)*

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Includes Health Post, Health Clinic, and Maternity Center

[3]Includes Chemist and Patent Medicine Vendors

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Contraceptive choice: Availability of modern contraception, by method

Percentage of service delivery points (SDPs) offering different types of contraceptive methods[1], by sector and method, among all SDPs

	Public SDPs						
	Contraceptive method						
SDP characteristic	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception
Number = 24							
Total	(66.7)	(66.7)	(12.5)	(37.5)	(50.0)	(0.0)	(25.0)
Facility type							
Hospital	(80.0)	(80.0)	(40.0)	(60.0)	(60.0)	(0.0)	(20.0)
Health center[2]	(63.2)	(63.2)	(5.3)	(31.6)	(47.4)	(0.0)	(26.3)
Pharmacy[3]	na	na	na	na	na	na	na
Other	na	na	na	na	na	na	na

	Private SDPs						
	Contraceptive method						
SDP characteristic	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception
Number = 20							
Total	(60.0)	(40.0)	(5.0)	(5.0)	(20.0)	(0.0)	(5.0)
Facility type							
Hospital	na	na	na	na	na	na	na
Health center[2]	*(100.0)*	*(100.0)*	*(100.0)*	*(100.0)*	*(0.0)*	*(0.0)*	*(0.0)*
Pharmacy[3]	(61.1)	(38.9)	(0.0)	(0.0)	(22.2)	(0.0)	(5.6)
Other	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Includes Health Post, Health Clinic, and Maternity Center

[3]Includes Chemist and Patent Medicine Vendors

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Contraceptive stock-outs, by method

Percentage of service delivery points (SDPs) stocked out of specific modern contraceptives[1] at time of survey or stocked out in the last 3 months, by sector and background characteristics, among SDPs that offer each method

	Public SDPs					
	Contraceptive method					
Background characteristic	Pill	Injectables	IUD	Implants	Male condom	Emergency contraception
Number of SDPs	16	16	3	9	12	6
Total	(12.5)	(31.3)	*(0.0)*	(44.4)	(33.3)	(33.3)
SDP type						
Hospital	*(0.0)*	*(25.0)*	*(0.0)*	*(33.3)*	*(33.3)*	*(0.0)*
Health center[2]	(16.7)	(33.3)	*(0.0)*	(50.0)	(33.3)	(40.0)
Pharmacy[3]	na	na	na	na	na	na
Other	na	na	na	na	na	na

	Private SDPs					
	Contraceptive method					
Background characteristic	Pill	Injectables	IUD	Implants	Male condom	Emergency contraception
Number of SDPS	12	8	1	1	4	1
Total	(16.7)	(25.0)	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*
SDP type						
Hospital	na	na	na	na	na	na
Health center[2]	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	na	na
Pharmacy[3]	(18.2)	(28.6)	na	na	*(0.0)*	*(0.0)*

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Includes Health Post, Health Clinic, and Maternity Center

[3]Includes Chemist and Patent Medicine Vendors

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Other	na	na	na	na	na	na

Number of family planning visits, by method

Total number of visits and new clients for contraceptive services, by method type and background characteristics, among advanced service delivery points (SDPs) that offer each method

	Contraceptive Methods														
	Sterilization		Injectable		IUD		Implants		Male condom		Pills		Emergency Contraception	
	Total women	Total men	Total	New	Total	New	Total	New	Total	New	Total	New	Total	New
Number of SDPs	0	0	17		4		10		12		17		6	
Total	na	na	407	243	13	6	163	129	12	7	166	91	4	3
Background characteristic														
SDP type														
Hospital	na	na	251	168	6	6	109	109	0	0	39	36	0	0
Health center[1]	na	na	156	75	7	0	54	20	12	7	127	55	4	3
Sector														
Public	na	na	387	238	6	6	145	126	12	7	136	81	4	3
Private	na	na	20	5	7	0	18	3	0	0	30	10	0	0

[1]Includes Health Post, Health Clinic, and Maternity Center