Background characteristics of respondents

Percent distribution of background characteristics of women age 15 to 49

Background characteristics	Urban	Total	Percentage of sample	Weighted number	Unweighted number
Age group					
15–19	100.0	100.0	13.8	196	193
20–24	100.0	100.0	13.0	183	184
25–29	100.0	100.0	17.8	250	260
30–34	100.0	100.0	18.8	264	263
35–39	100.0	100.0	17.9	251	259
40–44	100.0	100.0	11.5	162	165
45–49	100.0	100.0	7.1	100	105
Marital status					
Married	100.0	100.0	61.0	858	878
Living together	100.0	100.0	4.9	69	73
Divorced	100.0	100.0	2.9	41	40
Widowed	(100.0)	(100.0)	1.7	24	23
Never married	100.0	100.0	29.5	415	414
Parity					
None	100.0	100.0	34.1	479	476
1-2	100.0	100.0	29.8	419	433
3-4	100.0	100.0	27.2	383	396
5 or more	100.0	100.0	8.5	120	119
Education					
Never attended	100.0	100.0	2.4	33	32
Primary	100.0	100.0	14.1	199	203
Secondary	100.0	100.0	53.7	755	791
Higher	100.0	100.0	29.7	418	401
Wealth quintile					
Lowest	100.0	100.0	15.8	223	215
Lower	100.0	100.0	18.1	255	295
Middle	100.0	100.0	21.7	305	315
Higher	100.0	100.0	21.5	302	320
Highest	100.0	100.0	22.9	323	284
Total	100.0	100.0	100.0	1,407	1,429

^{() =} Estimate generated from less than 25 unweighted observations

Current use of contraception by background characteristics

Percentage of women ages 15 to 49 currently using a contraceptive method, by type and background characteristics

	Any me	ethod	Modern r	method ¹	Tradition	al method
Background characteristic	All women	Married women ²	All women	Married women ²	All women	Married women ²
Number	1,407	926	1,407	926	1,407	926
	28.0	34.9	21.0	26.4	6.8	8.3
Total	40.0	34.7	41.0	∠ ∪. +	0.0	0.3
Age group 15–19	3.2	(0,0)	3.2	(0,0)	0.0	(0,0)
		(0.0)		(0.0)		(0.0)
20–24	23.3	17.8	18.3	16.6	5.0	1.2
25–29	25.9	26.9	21.5	23.0	4.4	3.9
30–34	39.3	42.7	30.6	32.7	8.7	10.0
35–39	36.4	39.3	24.3	26.6	11.0	11.5
40–44	32.5	35.2	24.4	26.8	8.1	8.4
45–49	31.8	36.6	20.8	25.4	11.0	11.1
Marital status						
Married ²	34.9	na	26.4	na	8.3	na
Unmarried, sexually active	42.3	na	31.6	na	10.7	na
Parity						
0–1	15.1	16.8	12.0	14.1	3.1	2.7
2–3	35.0	36.2	26.7	27.8	8.3	8.4
4 or more	47.7	50.4	33.8	35.9	13.0	13.5
Education						
Never attended	14.4	(18.2)	14.4	(18.2)	0.0	(0.0)
Primary	22.7	25.0	15.3	16.8	7.4	8.1
Secondary	27.0	36.8	20.8	28.1	5.9	8.1
Higher	33.3	39.8	24.9	30.3	8.5	9.5
Wealth quintile						
Lowest	17.8	23.5	13.7	17.9	4.1	5.6
Lower	21.2	24.3	16.3	18.3	4.9	6.0
Middle	27.5	30.4	20.4	24.1	7.2	6.4
Higher	31.8	39.1	24.9	29.9	6.9	9.1
Highest	37.1	50.6	26.9	36.7	9.5	12.7

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na = not applicable

²Married includes married women and those who are living with a man

Contraceptive method mix by background characteristics

Percent distribution of current contraceptive users ages 15 to 49, by method type, marital status and background characteristics

						All users				
						N = 393				
Background	Sterili-			Inject-		Emergency		Other	Other	
characteristics	zation	Implants	IUD	ables	Pill	contraception	Condom	modern	traditional	Total
Total	2.4	4.0	7.5	13.8	13.6	5.5	27.1	3.0	23.0	100.0
Age group										
15–19	(0.0)	(0.0)	(0.0)	(0.0)	(20.7)	(12.2)	(67.1)	(0.0)	(0.0)	(100.0)
20–24	0.0	5.2	0.0	3.8	16.5	19.6	33.2	0.0	21.7	100.0
25–29	0.0	1.0	5.9	10.7	13.5	8.9	43.1	0.0	17.1	100.0
30–34	1.0	2.4	13.2	10.5	16.5	3.5	26.2	4.7	22.1	100.0
35–39	4.2	5.8	6.0	13.5	9.6	3.1	24.9	7.5	25.5	100.0
40–44	2.5	5.4	11.5	28.1	16.3	0.9	10.2	0.0	25.0	100.0
45–49	10.6	7.2	1.8	24.1	5.9	0.0	16.0	0.0	34.5	100.0
Marital status										
Married ¹	3.0	4.9	9.2	16.4	14.5	2.6	23.5	3.6	22.4	100.0
Unmarried, sexually	0.0	0.0	0.0	2.5	7.0	15.2	50.0	0.0	25.3	100.0
active										
Parity										
0–1	0.0	2.9	0.0	2.4	12.0	15.5	46.6	0.0	20.7	100.0
2–3	1.3	3.1	12.9	16.5	15.0	2.5	23.3	2.7	22.8	100.0
4 or more	5.6	6.0	6.9	19.1	13.1	1.7	17.1	5.5	25.1	100.0
Education										
Never attended	(0.0)	(0.0)	(0.0)	(61.1)	(0.0)	(18.7)	(20.2)	(0.0)	(0.0)	(100.0)
Primary	0.0	2.8	6.2	15.1	25.2	4.0	11.6	4.6	30.5	100.0
Secondary	0.5	3.4	4.4	16.8	15.0	6.0	28.8	3.4	21.7	100.0
Higher	6.2	5.4	12.8	7.2	8.2	4.8	29.9	2.0	23.4	100.0
Wealth quintile					-					
Lowest	0.0	3.2	3.3	11.4	19.1	9.0	31.1	2.6	20.2	100.0
Lower	0.0	5.4	3.7	17.7	11.1	9.9	25.9	3.0	23.2	100.0
Middle	1.2	5.3	3.9	12.3	14.9	4.5	31.8	0.6	25.5	100.0
Higher	0.0	2.8	5.8	16.5	18.9	4.7	28.0	1.4	21.8	100.0
Highest	7.2	3.7	14.5	11.6	7.7	3.8	22.4	6.1	23.2	100.0

¹Married includes married women and those who are living with a man

^{() =} Estimate generated from less than 25 unweighted observations


^{() =} Estimate generated from less than 5 unweighted observations

					M	arried users				
						N = 323				
Background	Sterili-			Inject-		Emergency		Other	Other	
characteristics	zation	Implants	IUD	ables	Pill	contraception	Condom	modern	traditional	Total
Total	3.0	4.9	9.2	16.4	14.5	2.6	23.5	3.6	22.4	100.0
Age group										
15–19	na	na	na	na	na	na	na	na	na	na
20–24	0.0	18.0	0.0	9.2	41.4	0.0	24.5	0.0	7.0	100.0
25–29	0.0	1.3	8.0	13.5	18.4	7.1	37.2	0.0	14.5	100.0
30–34	1.1	2.6	14.3	11.3	15.1	2.8	24.3	5.1	23.3	100.0
35–39	4.5	6.1	6.4	14.4	9.3	2.1	25.1	8.1	24.2	100.0
40–44	2.5	5.5	11.7	28.6	16.6	1.0	10.4	0.0	23.7	100.0
45–49	11.3	7.7	1.9	25.5	6.2	0.0	17.0	0.0	30.4	100.0
Marital status										
Married	na	na	na	na	na	na	na	na	na	na
Unmarried, sexually	na	na	na	na	na	na	na	na	na	na
active										
Parity										
0–1	0.0	7.1	0.0	4.7	19.4	9.8	43.1	0.0	16.0	100.0
2–3	1.4	3.2	13.4	16.8	14.5	1.4	24.2	2.8	22.3	100.0
4 or more	5.7	6.1	7.1	19.6	12.9	1.7	16.6	5.7	24.4	100.0
Education										
Never attended	(0.0)	(0.0)	(0.0)	(61.1)	(0.0)	(18.7)	(20.2)	(0.0)	(0.0)	(100.0)
Primary	0.0	3.0	6.6	16.1	24.9	2.1	12.4	4.9	30.1	100.0
Secondary	0.6	4.3	5.4	20.2	16.8	2.7	24.2	4.2	21.7	100.0
Higher	7.7	6.8	16.1	9.0	7.7	1.9	26.9	2.6	21.3	100.0
Wealth quintile										
Lowest	0.0	3.9	4.1	14.0	18.5	5.9	29.8	3.2	20.8	100.0
Lower	0.0	7.3	5.0	22.7	12.9	8.1	15.3	4.1	24.6	100.0
Middle	1.6	7.2	5.4	16.1	18.2	0.0	30.6	0.8	20.1	100.0
Higher	0.0	3.3	6.9	19.4	20.7	3.9	20.8	1.7	23.4	100.0
Highest	8.0	4.1	16.1	12.8	7.0	0.0	22.7	6.7	22.5	100.0

^{() =} Estimate generated from less than 25 unweighted observations

^{() =} Estimate generated from less than 5 unweighted observations

Current Method Mix Among **All and Married** Contraceptive Users (Age 15-49)



Unmet need for family planning

Percentage of women age 15 to 49 with unmet need for family planning, by marital status and background characteristics

Background characteristics		All women			Married women ¹	
Number		1,407			926	
	For spacing	For limiting	Total	For spacing	For limiting	Total
Total	9.9	7.9	17.8	12.4	11.3	23.7
Age group						
15–19	3.8	1.4	5.3	(28.6)	(11.4)	(40.0)
20–24	15.2	1.4	16.5	22.8	3.6	26.4
25–29	15.6	2.3	17.9	20.1	3.2	23.4
30–34	13.4	6.2	19.6	14.8	6.1	20.9
35–39	6.4	18.1	24.5	6.9	19.9	26.8
40–44	7.4	17.0	24.5	8.2	18.8	27.0
45–49	1.0	11.4	12.4	1.2	12.8	14.1
Marital status						
Married	12.4	11.3	23.7	na	na	na
Unmarried, sexually	24.2	5.9	30.1	na	na	na
active						
Parity						
0–1	8.2	1.2	9.3	13.2	1.3	14.6
2–3	14.9	9.5	24.4	15.7	10.2	25.9
4 or more	5.4	21.2	26.6	5.9	21.9	27.8
Education						
Never attended	11.5	12.3	23.8	(11.2)	(15.5)	(26.7)
Primary	11.9	12.5	24.4	13.2	13.0	26.2
Secondary	9.8	8.0	17.8	13.8	12.4	26.2
Higher	8.8	5.4	14.2	9.6	8.1	17.7
Wealth quintile						
Lowest	10.6	10.1	20.7	13.6	12.2	25.8
Lower	15.6	10.1	25.7	19.8	14.5	34.3
Middle	8.8	6.0	14.8	12.2	9.0	21.3
Higher	10.3	8.8	19.1	12.2	12.7	24.9
Highest	5.5	5.9	11.3	6.4	8.9	15.3

^{() =} Estimate generated from less than 25 unweighted observations

Need and demand for family planning

Percentage of women age 15 to 49 with met need, unmet need, and total demand for family planning, and percentage of demand for family planning satisfied with modern methods, by marital status and background characteristics

Background characteristics			All wome	n			M	arried wo	men ²	
Number			1,407					926		
	Any Method	Modern method ¹	Unmet need	Total demand	Demand satisfied by modern method	Any Method	Modern method ¹	Unmet need	Total demand	Demand satisfied by modern method
Total	28.0	21.0	17.8	45.6	46.1	34.9	26.4	23.7	58.3	45.2
Age group										
15–19	3.2	3.2	5.3	8.5	37.9	(0.0)	(0.0)	(40.0)	(40.0)	(0.0)
20–24	23.3	18.3	16.5	39.9	45.8	17.8	16.6	26.4	44.2	37.5
25–29	25.9	21.5	17.9	43.8	49.0	26.9	23.0	23.4	50.3	45.8
30–34	39.3	30.6	19.6	58.9	52.0	42.7	32.7	20.9	63.5	51.5
35–39	36.4	24.3	24.5	59.9	40.6	39.3	26.6	26.8	64.9	41.0
40–44	32.5	24.4	24.5	56.9	42.8	35.2	26.8	27.0	62.2	43.2
45–49	31.8	20.8	12.4	44.1	47.1	36.6	25.4	14.1	50.6	50.2
Marital status										
Married ²	34.9	26.4	23.7	58.3	45.2	na	na	na	na	na
Unmarried,	42.3	31.6	30.1	72.4	43.6	na	na	na	na	na
sexually active										
Parity										
0–1	15.1	12.0	9.3	24.4	49.0	16.8	14.1	14.6	31.4	45.0
2–3	35.0	26.7	24.4	59.4	45.0	36.2	27.8	25.9	62.2	44.8
4 or more	47.7	33.8	26.6	73.4	46.0	50.4	35.9	27.8	77.2	46.5
Education										
Never attended	14.4	14.4	23.8	38.2	37.7	(18.2)	(18.2)	(26.7)	(44.9)	(40.5)
Primary	22.7	15.3	24.4	47.1	32.4	25.0	16.8	26.2	51.1	32.9
Secondary	27.0	20.8	17.8	44.5	46.7	36.8	28.1	26.2	62.4	45.1
Higher	33.3	24.9	14.2	47.5	52.3	39.8	30.3	17.7	57.5	52.7
Wealth quintile										
Lowest	17.8	13.7	20.7	38.5	35.6	23.5	17.9	25.8	49.3	36.3
Lower	21.2	16.3	25.7	46.9	34.7	24.3	18.3	34.3	58.5	31.2
Middle	27.5	20.4	14.8	42.3	48.1	30.4	24.1	21.3	51.7	46.5
Higher	31.8	24.9	19.1	50.9	48.9	39.1	29.9	24.9	63.9	46.8
Highest	37.1	26.9	11.3	47.7	56.3	50.6	36.7	15.3	64.7	56.7

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na = not applicable

²Married includes married women and those who are living with a man

Pregnancy intentions

Unintended births among women age 15 to 49 with one or more births in the previous 5 years or who are currently pregnant: Percent distribution reporting last or current pregnancy was wanted then, wanted later or not wanted at all, by marital status and background characteristics

Background characteristics	All w	omen with	1 or more b	oirths	Married women with 1 or more births			
Number		6	27			586		
	Wanted then	Wanted later	Wanted no more	Total	Wanted then	Wanted later	Wanted no more	Total
Total	72.6	14.2	13.3	100.0	74.9	13.1	11.9	100.0
Age group								
15–19	(43.3)	(19.8)	(37.0)	(100.0)	(70.6)	(10.3)	(19.1)	(100.0)
20–24	66.1	20.0	13.9	100.0	78.3	16.7	5.0	100.0
25–29	77.5	18.4	4.1	100.0	78.0	17.7	4.4	100.0
30–34	75.6	14.1	10.3	100.0	77.4	13.9	8.7	100.0
35–39	69.5	9.5	21.0	100.0	69.4	9.4	21.1	100.0
40–44	70.4	5.3	24.4	100.0	71.5	2.8	25.7	100.0
45–49	(67.9)	(22.3)	(9.8)	(100.0)	(65.4)	(24.0)	(10.6)	(100.0)
Marital status								
Married ¹	74.9	13.1	11.9	100.0	na	na	na	na
Unmarried, sexually active	(21.4)	(39.1)	(39.5)	(100.0)	na	na	na	na
Parity								
0–1	73.5	17.5	9.0	100.0	82.5	14.2	3.3	100.0
2–3	78.6	12.6	8.9	100.0	78.8	12.8	8.4	100.0
4 or more	60.1	13.7	26.2	100.0	60.5	12.8	26.7	100.0
Education								
Never attended	(82.2)	(10.9)	(6.9)	(100.0)	(80.9)	(11.7)	(7.4)	(100.0)
Primary	71.6	8.0	20.3	100.0	73.9	8.8	17.3	100.0
Secondary	72.0	14.1	13.8	100.0	75.5	12.4	12.1	100.0
Higher	73.3	17.6	9.1	100.0	74.0	16.6	9.4	100.0
Wealth quintile								
Lowest	66.7	16.1	17.2	100.0	74.0	14.4	11.7	100.0
Lower	80.6	7.0	12.4	100.0	85.2	3.9	10.9	100.0
Middle	73.7	16.7	9.7	100.0	74.8	16.8	8.5	100.0
Higher	74.9	13.6	11.5	100.0	75.1	13.4	11.5	100.0
Highest	66.3	16.9	16.7	100.0	67.1	15.8	17.1	100.0

¹Married includes married women and those who are living with a man

na = not applicable

Contraceptive choice

Percent distribution of women age 15 to 49 who are currently using modern contraception¹ reporting on who decided on the contraceptive method, by background characteristics

Background characteristics	Self	Jointly with partner or provider	Provider alone/ partner alone/other	Total
Number	131	214	41	386
Total	33.9	55.4	10.7	100.0
Age group				
15–19	(32.9)	(25.7)	(41.4)	(100.0)
20–24	44.0	49.2	6.8	100.0
25–29	31.7	62.1	6.2	100.0
30–34	33.9	56.0	10.1	100.0
35–39	33.7	55.2	11.1	100.0
40–44	34.8	52.6	12.6	100.0
45–49	23.6	59.8	16.6	100.0
Marital status				
Married ²	30.7	58.8	10.5	100.0
Unmarried, sexually active	39.9	40.7	19.4	100.0
Parity				
0–1	36.7	52.8	10.4	100.0
2–3	26.5	61.8	11.7	100.0
4 or more	40.2	50.1	9.7	100.0
Education				
Never attended	(32.5)	(47.3)	(20.2)	(100.0)
Primary	47.2	47.0	5.7	100.0
Secondary	32.0	53.5	14.5	100.0
Higher	32.3	61.3	6.4	100.0
Wealth quintile				
Lowest	52.3	32.0	15.7	100.0
Lower	48.3	44.8	6.9	100.0
Middle	31.3	58.8	9.9	100.0
Higher	25.4	61.2	13.4	100.0
Highest	29.7	61.3	9.1	100.0

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

^{() =} Estimate generated from less than 25 unweighted observations

^{() =} Estimate generated from less than 5 unweighted observations

Payment for family planning

Percentage of women age 15 to 49 currently using a modern contraceptive method¹ who paid fees for services, by marital status and background characteristics

Background characteristics	All modern contraceptive users ¹	Married ² modern contraceptive users ¹
Number	393	323
Total	39.2	41.4
Age group		
15–19	(12.2)	na
20–24	39.8	(63.5)
25–29	50.9	59.6
30–34	48.7	49.1
35–39	31.7	31.2
40–44	32.6	33.1
45–49	21.4	22.8
Marital status		
Married ²	41.4	na
Unmarried, sexually active	35.2	na
Parity		
0–1	38.7	58.7
2–3	44.7	44.0
4 or more	33.2	33.2
Education		
Never attended	(38.9)	(38.9)
Primary	38.8	39.2
Secondary	43.9	47.5
Higher	32.4	33.4
Wealth quintile		
Lowest	37.8	38.6
Lower	44.4	48.0
Middle	44.0	50.7
Higher	44.6	48.3
Highest	29.7	29.5

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na = not applicable

- () = Estimate generated from less than 25 unweighted observations
- () = Estimate generated from less than 5 unweighted observations

²Married includes married women and those who are living with a man

Method Information Index Components

Percentage of current modern¹ contraceptive users who reported whether provider informed them about other methods, side effects and, if informed of side effects, what to do, by marital status and background characteristics²

Background characteristics		ned about methods		ned about effects	Told what to do if experienced side effects		
	All users	Married users ³	All users	Married users ³	All users	Married users ³	
Number	393	323	308	250	113	103	
Total	43.0	49.6	36.7	41.2	82.3	83.8	
Age group							
15–19	(0.0)	na	(0.0)	na	na	na	
20–24	16.6	(47.9)	34.0	(50.9)	(67.5)	(57.1)	
25–29	26.3	30.0	23.1	27.4	(78.5)	(80.0)	
30–34	49.3	50.7	34.4	34.4	72.4	75.8	
35–39	53.3	55.3	30.9	33.3	(96.6)	(96.6)	
40–44	53.9	54.8	65.3	66.6	84.2	84.2	
45–49	53.0	52.9	(57.0)	(59.2)	(95.9)	(95.9)	
Marital status							
Married ³	49.6	na	41.2	na	83.8	na	
Unmarried, sexually	12.6	na	16.3	na	(90.9)	na	
active							
Parity							
0–1	16.7	30.7	19.4	24.5	71.6	60.0	
2–3	50.3	50.1	42.7	42.6	81.1	84.4	
4 or more	54.2	55.0	43.9	45.3	87.7	87.7	
Education							
Never attended	(61.1)	(61.1)	(81.3)	(81.3)	(100.0)	(100.0)	
Primary	62.5	64.2	30.7	32.6	(75.9)	(75.9)	
Secondary	37.2	43.1	35.2	40.2	81.1	83.0	
Higher	44.7	53.3	38.8	43.3	83.8	85.3	
Wealth quintile							
Lowest	48.8	54.8	42.2	(45.9)	(55.6)	(63.0)	
Lower	47.2	60.0	36.9	45.5	(93.0)	(95.6)	
Middle	32.1	38.2	23.7	29.4	(82.1)	(81.0)	
Higher	41.9	49.3	38.6	41.6	79.3	81.3	
Highest	47.7	51.0	42.5	45.0	89.8	89.3	

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na = not applicable

- () = Estimate generated from less than 25 unweighted observations
- () = Estimate generated from less than 5 unweighted observations

²"Informed about other methods" asked of all current contraceptive users, "informed about side effects" asked of current modern contraceptive users, and "told what to do if experienced side effects" asked of current modern contraceptive users who indicated they were told about side effects.

³Married includes married women and those who are living with a man

Perceived quality of care for family planning

Percentage of current modern¹ contraceptive users who would return, would refer a friend/relative, or would return and refer friend/relative to provider, by background characteristics

Background characteristics	Percent who would return to provider	Percent who would refer relative/friend	Percent who would return and refer friend/relative
Number	306	306	306
Total	73.6	73.5	68.8
Age group			
15–19	(15.9)	(15.9)	(15.9)
20–24	51.9	47.4	47.4
25–29	77.8	73.1	73.1
30–34	78.5	79.2	74.1
35–39	83.8	81.9	76.4
40–44	81.4	82.8	76.2
45–49	(53.8)	(71.2)	(53.8)
Marital status			
Married ²	79.2	79.8	74.5
Unmarried, sexually active	61.1	57.8	57.8
Parity			
0–1	58.2	55.1	53.6
2–3	79.9	79.8	75.6
4 or more	78.8	81.2	73.3
Education			
Never attended	(52.7)	(52.7)	(52.7)
Primary	84.8	81.9	81.9
Secondary	67.4	69.2	63.7
Higher	80.9	78.7	73.7
Wealth quintile			
Lowest	58.2	63.2	54.9
Lower	77.2	73.9	73.9
Middle	70.9	66.5	66.5
Higher	70.9	73.9	66.2
Highest	81.5	81.6	75.2

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na = not applicable

- () = Estimate generated from less than 25 unweighted observations
- () = Estimate generated from less than 5 unweighted observations

²Married includes married women and those who are living with a man

Knowledge of family planning

Percentage of women age 15 to 49 who reported receiving family planning information from a health worker in the past 12 months, by marital status and background characteristics

Background characteristics		ealth worker who amily planning		ility and provider family planning	Either visited by health worker or visited facility and discussed family planning		
Number	All women 1,407	Married women ¹ 926	All women 1,407	Married women ¹ 926	All women 1,407	Married women ¹ 926	
Total	9.6	12.2	24.3	34.4	28.9	39.6	
Age group							
15–19	4.7	(8.1)	3.0	(14.3)	7.3	(14.3)	
20–24	10.0	21.9	20.5	48.9	23.7	52.9	
25–29	9.5	12.2	30.8	41.9	35.4	47.6	
30–34	15.1	15.8	37.0	41.4	42.4	46.5	
35–39	9.8	10.5	31.2	34.7	36.0	39.8	
40–44	9.7	9.3	20.3	19.3	26.7	26.0	
45–49	3.8	4.7	12.2	14.9	14.6	17.8	
Marital status							
Married ¹	12.2	na	34.4	na	39.6	na	
Unmarried,	4.8	na	4.9	na	7.1	na	
sexually active							
Parity							
0–1	5.0	7.1	13.4	30.4	16.6	33.3	
2–3	15.5	15.7	35.8	37.7	42.9	44.6	
4 or more	10.8	11.1	32.2	33.8	35.7	37.6	
Education							
Never attended	8.5	(10.8)	23.8	(30.1)	28.4	(35.9)	
Primary	11.3	11.7	23.2	26.1	30.7	34.4	
Secondary	11.0	15.6	24.5	38.2	29.5	44.1	
Higher	6.5	7.4	24.5	34.0	27.1	36.1	
Wealth quintile							
Lowest	11.5	16.5	16.7	23.2	23.7	33.1	
Lower	12.5	17.4	24.3	35.5	29.5	42.1	
Middle	11.0	15.1	29.9	42.8	34.0	48.0	
Higher	7.9	9.0	29.3	40.2	32.2	42.5	
Highest	6.5	6.0	19.6	27.3	24.1	31.0	

¹Married includes married women and those who are living with a man

na = not applicable

List of indicators for sampling errors, PMA2015/Lagos-Round 2

Variable	Estimation	Base population
Currently using a modern method	Proportion	All women ages 15–49
Currently using a modern method	Troportion	Married women ages 15–49
Currently using a traditional method	Proportion	All women ages 15–49
Currently using a traditional method	Troportion	Married women ages 15–49
Currently using any contraceptive method	Proportion	All women ages 15–49
Currently using any contraceptive method	Troportion	Married women ages 15–49
Currently using injectables	Proportion	All women ages 15–49
Currently using injectables	Troportion	Married women ages 15–49
Cumontly using male condoms	Proportion	All women ages 15–49
Currently using male condoms	Troportion	Married women ages 15–49
Commental variage invalents	Proportion	All women ages 15–49
Currently using implants	Troportion	Married women ages 15–49
Chase method by self or identity in most 12 months	Proportion	All women ages 15–49
Chose method by self or jointly in past 12 months	Troportion	Married women ages 15–49
Doid food for family planning convices in past 12 months	Proportion	All women ages 15–49
Paid fees for family planning services in past 12 months	Troportion	Married women ages 15–49
Informed by mayiden about other methods	Proportion	All women ages 15–49
Informed by provider about other methods	Troportion	Married women ages 15–49
Informed by mayidar shout side offeets	Proportion	All women ages 15–49
Informed by provider about side effects	Troportion	Married women ages 15–49
Satisfied with provider: Would return and refer	Duo no anti o ::	All women ages 15–49
friend/relative to provider	Proportion	Married women ages 15–49
_		A11 15 40
Visited by health worker who talked about family planning	Proportion	All women ages 15–49
information in past 12 months	_	Married women ages 15–49

Characteristics of service delivery points

Number and percentage of service delivery points (SDPs) by background characteristics

Background characteristics	Number	Percentage
Total	123	100.0
SDP type		
Hospital	25	20.3
Health center ¹	56	45.5
Pharmacy ²	40	32.5
Sector		
Public	67	54.5
Private	56	45.5
Offers family planning		
No	9	7.3
Yes	114	92.7

¹Includes Health Post, Health Clinic, and Maternity Center

²Includes Chemist and Patent Medicine Vendors

Charging fees for family planning

Percentage of service delivery points (SDPs) that charge fees for family planning services, by sector and background characteristics, among SDPs offering family planning

Background characteristic	Sec	ctor
	Public	Private
Number of SDPs	65	49
Total	9.2	61.2
SDP type		
Hospital	(21.4)	(88.9)
Health center ¹	5.9	(100.0)
Pharmacy ²	na	52.8

¹Includes Health Post, Health Clinic, and Maternity Center

- () = Estimate generated from less than 25 unweighted observations
- () = Estimate generated from less than 5 unweighted observations

²Includes Chemist and Patent Medicine Vendors

Contraceptive choice: Availability of at least 3 or 5 modern contraceptive methods

Percentage of public and private service delivery points (SDPs) offering at least 3 or 5 modern contraceptive methods¹, by sector and background characteristics, among all SDPs

Background characteristic	3 or more	e methods	5 or more methods			
	Public	Private	Public	Private		
Number of SPDs	67	56	67	56		
Total	95.5	51.8	92.5	16.1		
SDP type						
Hospital	(100.0)	(63.6)	(100.0)	(36.4)		
Health Center ²	94.3	(66.7)	90.6	(66.7)		
Pharmacy ³	na	47.5	na	7.5		

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na = not applicable

() = Estimate generated from less than 25 unweighted observations

²Includes Health Post, Health Clinic, and Maternity Center

³Includes Chemist and Patent Medicine Vendors

Contraceptive Choice: Availability of modern contraception, by method

Percentage of service delivery points (SDPs) offering different types of contraceptive methods¹, by sector and method, among all SDPs

	Public SDPs									
	Contraceptive method									
Background characteristic	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception			
Number of SDPs = 67										
Total	97.0	97.0	85.1	79.1	97.0	11.9	37.3			
SDP type										
Hospital	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(57.1)	(71.4)			
Health center ²	96.2	96.2	81.1	73.6	96.2	0.0	28.3			
Pharmacy ³	na	na	na	na	na	na	na			

	Private SDPs										
	Contraceptive method										
Background characteristic	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception				
Number of SDPs = 56											
Total	69.6	28.6	19.6	12.5	75.0	7.1	46.4				
SDP type											
Hospital	(63.6)	(72.7)	(72.7)	(36.4)	(45.5)	(36.4)	(36.4)				
Health center ²	(66.7)	(66.7)	(66.7)	(66.7)	(66.7)	(0.0)	(33.3)				
Pharmacy ³	70.0	12.5	0.0	0.0	82.5	0.0	50.0				

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Includes Health Post, Health Clinic, and Maternity Center

³Includes Chemist and Patent Medicine Vendors

^{() =} Estimate generated from less than 25 unweighted observations

^{() =} Estimate generated from less than 5 unweighted observations

Contraceptive stock-outs, by method

Percentage of service delivery points (SDPs) stocked out of specific modern contraceptives¹ at time of survey or stocked out in the last 3 months, by sector and background characteristics, among SDPs that offer each method

		Public SDPs								
	Contraceptive method									
Background characteristic	Male Emergency Pill Injectables IUD Implants condom contraception									
Number of SPDs	65	65	57	53	65	25				
Total	3.1	6.2	1.8	13.2	6.2	12.0				
SDP type										
Hospital	(0.0)	(14.3)	(7.1)	(14.3)	(7.1)	(20.0)				
Health center ¹	3.9	3.9	0.0	12.8	5.9	(6.7)				
Pharmacy ²	na	na	na	na	na	na				

	Private SDPs									
	Contraceptive method									
Background characteristic	Male Emergency Pill Injectables IUD Implants condom contraception									
Number of SDPs	39	16	11	7	42	26				
Total	23.1	(12.5)	(18.2)	(28.6)	19.0	23.1				
SDP type										
Hospital	(0.0)	(0.0)	(12.5)	(25.0)	(0.0)	(0.0)				
Health center ¹	(0.0)	(0.0)	(0.0)	(50.0)	(0.0)	(0.0)				
Pharmacy ²	28.6	(20.0)	na	na	18.2	30.0				

¹Includes Health Post, Health Clinic, and Maternity Center

²Includes Chemist and Patent Medicine Vendors

^{() =} Estimate generated from less than 25 unweighted observations

^{() =} Estimate generated from less than 5 unweighted observations

Number of family planning visits, by method

Total number of visits and new clients contraceptive services, by method type and background characteristics, among advanced service delivery points (SDPs) that offer each method

	Contraceptive Methods														
	Sterilization Injectable		IUD Implants		Male condom		Pills		Emergency Contraception						
	Total women	Total men	Total	New	Total	New	Total	New	Total	New	Total	New	Total	New	
Number of SPDs	12	11		76		68		60		74		76		31	
Total	5	0	1,809	382	586	225	525	296	3,724	1,504	743	347	25	13	
Background characteristic															
SDP type															
Hospital	5	0	511	125	369	139	204	111	599	297	330	123	22	10	
Health center ¹	na	na	1,292	256	214	85	321	185	3,120	1,207	413	224	0	0	
Sector															
Public	5	0	1,743	369	512	179	513	291	3,659	1,484	676	336	12	8	
Private	0	0	66	13	74	46	12	5	65	20	67	11	13	5	

¹Includes Health Post, Health Clinic, and Maternity Center