

List of indicators for sampling errors, PMA2016/Kano

| Variable | Estimation | Base population |
|---|------------|--|
| Currently using a modern method | Proportion | All women ages 15–49 Married women ages 15–49 |
| Currently using a traditional method | Proportion | All women ages 15–49 Married women ages 15–49 |
| Currently using any contraceptive method | Proportion | All women ages 15–49 Married women ages 15–49 |
| Currently using injectables | Proportion | All women ages 15–49 Married women ages 15–49 |
| Currently using male condoms | Proportion | All women ages 15–49 Married women ages 15–49 |
| Currently using implants | Proportion | All women ages 15–49 Married women ages 15–49 |
| Chose method by self or jointly in past 12 months | Proportion | All women ages 15–49 Married women ages 15–49 |
| Paid fees for family planning services in past 12 months | Proportion | All women ages 15–49 Married women ages 15–49 |
| Informed by provider about other methods | Proportion | All women ages 15–49 Married women ages 15–49 |
| Informed by provider about side effects | Proportion | All women ages 15–49 Married women ages 15–49 |
| Satisfied with provider: Would return and refer friend/relative to provider | Proportion | All women ages 15–49 Married women ages 15–49 |
| Visited by health worker who talked about family planning information in past 12 months | Proportion | All women ages 15–49 Married women ages 15–49 |