

Background characteristics of respondents**Percent distribution of background characteristics of women age 15 to 49**

Background characteristics	Urban	Rural	Total	Percentage of sample	Weighted number	Unweighted number
Age group						
15–19	71.2	28.8	100.0	18.3	239	249
20–24	74.3	25.7	100.0	17.2	225	222
25–29	75.8	24.2	100.0	18.2	238	232
30–34	75.1	24.9	100.0	16.2	212	195
35–39	73.5	26.5	100.0	13.4	175	177
40–44	67.0	33.0	100.0	9.1	119	124
45–49	69.0	31.0	100.0	7.7	101	110
Marital status						
Married	71.2	28.8	100.0	50.6	662	662
Living together	(76.9)	(23.1)	(100.0)	0.9	12	10
Divorced	(65.3)	(34.7)	(100.0)	1.7	22	23
Widowed	70.5	29.5	100.0	4.2	54	57
Never married	75.5	24.5	100.0	42.6	558	557
Parity						
None	75.6	24.4	100.0	47.3	620	612
1-2	71.9	28.1	100.0	18.9	247	239
3-4	70.0	30.0	100.0	17.9	235	238
5 or more	69.2	30.8	100.0	15.5	203	214
Education						
Never attended	(59.8)	(40.2)	(100.0)	0.5	6	8
Primary	65.4	34.6	100.0	7.9	104	122
Secondary	69.8	30.2	100.0	61.5	806	803
Higher	81.5	18.5	100.0	29.9	392	375
Wealth quintile						
Lowest	47.9	52.1	100.0	16.0	209	278
Lower	67.0	33.0	100.0	20.4	267	276
Middle	75.2	24.8	100.0	20.3	265	257
Higher	78.8	21.2	100.0	21.2	277	235
Highest	88.8	11.2	100.0	22.2	290	263
Total	72.9	27.1	100.0	100.0	1,309	1,309

() = Estimate generated from less than 25 unweighted observations

Current use of contraception by background characteristics**Percentage of women ages 15 to 49 currently using a contraceptive method, by type and background characteristics**

Background characteristic	Any Method		Modern method ¹		Traditional method	
	All women	Married women ²	All women	Married women ²	All women	Married women ²
Number	1,309	674	1,309	674	1,309	674
Total	25.1	33.5	15.6	17.5	9.5	15.9
Age group						
15–19	6.9	(17.4)	6.2	(17.4)	0.7	(0.0)
20–24	24.9	26.6	21.3	20.0	3.5	6.6
25–29	26.7	26.3	17.6	14.5	9.2	11.7
30–34	33.2	38.8	16.6	17.5	16.5	21.3
35–39	38.8	41.7	20.4	20.6	18.4	21.2
40–44	24.2	28.5	15.1	17.8	9.1	10.7
45–49	25.5	34.7	11.1	14.4	14.4	20.3
Marital status						
Married ²	33.5	na	17.5	na	15.9	na
Unmarried, sexually active	36.7	na	31.2	na	5.5	na
Parity						
0–1	15.4	13.9	12.3	9.0	3.1	4.9
2–3	29.3	29.7	17.6	16.8	11.8	12.9
4 or more	44.7	48.8	22.2	23.5	22.6	25.3
Residence						
Urban	26.4	35.1	16.2	17.3	10.2	17.7
Rural	21.7	29.6	14.2	18.1	7.5	11.5
Education						
Never attended	(7.0)	(0.0)	(0.0)	(0.0)	(7.0)	(0.0)
Primary	25.1	33.9	11.9	16.4	13.3	17.5
Secondary	22.6	33.2	13.9	17.4	8.6	15.8
Higher	30.8	34.6	20.5	18.5	10.3	16.0
Wealth quintile						
Lowest	19.2	28.2	13.5	18.8	5.7	9.4
Lower	20.2	28.6	11.2	11.3	9.0	17.3
Middle	26.8	32.6	18.6	20.3	8.3	12.3
Higher	31.3	37.6	18.1	17.6	13.2	19.9
Highest	26.5	37.4	16.3	19.2	10.2	18.2

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

Contraceptive method mix by background characteristics

Percent distribution of current contraceptive users ages 15 to 49, by method type, marital status and background characteristics

All users										
N = 329										
Background characteristics	Sterilization	Implants	IUD	Injectables	Pill	Emergency contraception	Condom	Other modern	Other traditional	Total
Total	1.1	4.4	2.8	3.7	11.7	1.5	36.3	2.4	36.2	100.0
Age group										
15–19	(0.0)	(0.0)	(0.0)	(0.0)	(21.3)	(0.0)	(68.7)	(0.0)	(10.0)	(100.0)
20–24	0.0	2.9	0.0	0.0	24.5	3.6	53.9	0.9	14.2	100.0
25–29	4.2	0.9	0.0	6.8	4.3	1.3	48.1	0.0	34.3	100.0
30–34	0.0	11.6	1.5	1.0	7.6	0.5	26.1	5.5	46.1	100.0
35–39	0.7	5.3	5.5	3.1	11.7	0.0	26.4	3.5	43.8	100.0
40–44	1.5	0.0	10.4	13.4	15.3	3.5	18.4	0.0	37.5	100.0
45–49	(0.0)	(2.0)	(6.1)	(4.4)	(2.8)	(2.2)	(22.1)	(3.6)	(56.6)	(100.0)
Marital status										
Married ¹	0.4	5.2	4.1	4.8	8.4	0.7	27.8	3.2	45.4	100.0
Unmarried, sexually active	0.0	0.0	0.0	2.5	21.4	1.8	59.4	0.0	15.0	100.0
Parity										
0–1	2.4	0.3	0.0	1.2	17.8	2.1	55.7	0.5	20.1	100.0
2–3	0.0	5.6	3.0	1.2	7.2	1.2	41.7	7.0	33.2	100.0
4 or more	0.6	7.1	5.0	6.9	9.1	1.1	18.2	1.5	50.4	100.0
Residence										
Urban	1.1	4.7	2.9	2.7	10.7	1.3	36.8	3.1	36.7	100.0
Rural	1.1	3.4	2.7	6.9	14.9	1.8	34.5	0.0	34.6	100.0
Education										
Never attended	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(100.0)	(100.0)
Primary	3.3	2.3	0.0	17.6	5.6	0.0	18.4	0.0	52.8	100.0
Secondary	0.0	5.3	2.3	1.8	14.2	1.0	36.8	1.7	36.8	100.0
Higher	2.2	3.5	4.3	3.6	9.2	2.5	39.5	3.8	31.4	100.0
Wealth quintile								0.0		
Lowest	2.1	2.6	0.0	11.7	16.2	3.5	34.1	0.0	29.7	100.0
Lower	0.0	3.0	0.0	0.0	6.9	1.8	43.7	0.0	44.6	100.0
Middle	0.0	10.8	0.0	1.0	11.4	0.5	44.6	0.7	30.9	100.0
Higher	0.0	2.3	5.5	1.3	10.0	1.2	36.4	5.7	37.6	100.0
Highest	3.5	2.8	6.0	7.3	14.8	1.3	24.3	2.9	37.1	100.0

¹Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Married users										
N = 226										
Background characteristics	Sterilization	Implants	IUD	Injectables	Pill	Emergency contraception	Condom	Other modern	Other traditional	Total
Total	0.4	5.2	4.1	4.8	8.4	0.7	27.8	3.2	45.4	100.0
Age group										
15–19	(0.0)	(0.0)	(0.0)	(0.0)	(19.1)	(0.0)	(80.9)	(0.0)	(0.0)	(100.0)
20–24	(0.0)	(8.9)	(0.0)	(0.0)	(24.6)	(0.0)	(41.8)	(0.0)	(24.7)	(100.0)
25–29	0.0	1.9	0.0	9.3	0.0	0.0	44.2	0.0	44.6	100.0
30–34	0.0	11.0	1.7	1.2	7.3	0.0	21.8	6.1	50.9	100.0
35–39	0.8	3.6	6.5	3.7	7.6	0.0	27.0	4.2	46.5	100.0
40–44	1.5	0.0	10.4	13.4	15.3	3.5	18.4	0.0	37.5	100.0
45–49	(0.0)	(2.2)	(6.5)	(4.7)	(3.0)	(2.4)	(18.9)	(3.8)	(58.5)	(100.0)
Marital status										
Married ¹	na	na	na	na	na	na	na	na	na	na
Unmarried, sexually active	na	na	na	na	na	na	na	na	na	na
Parity										
0–1	(0.0)	(1.5)	(0.0)	(0.0)	(8.9)	(0.0)	(54.7)	(0.0)	(35.0)	(100.0)
2–3	0.0	5.6	3.3	1.3	7.9	0.0	38.5	7.7	35.6	100.0
4 or more	0.6	5.7	5.3	7.3	8.5	1.2	18.0	1.6	51.9	100.0
Residence										
Urban	0.0	6.2	4.3	3.3	5.3	0.6	28.5	4.3	47.6	100.0
Rural	1.5	2.5	3.7	9.2	17.6	1.0	25.6	0.0	38.9	100.0
Education										
Never attended	na	na	na	na	na	na	na	na	na	na
Primary	(3.9)	(2.7)	(0.0)	(20.6)	(6.6)	(0.0)	(14.7)	(0.0)	(51.6)	(100.0)
Secondary	0.0	5.4	3.3	2.5	11.5	0.0	29.7	2.0	45.6	100.0
Higher	0.0	5.7	6.9	3.9	3.6	2.1	28.5	6.1	43.2	100.0
Wealth quintile										
Lowest	3.1	3.7	0.0	16.9	18.5	2.1	22.3	0.0	33.4	100.0
Lower	0.0	4.6	0.0	0.0	10.5	0.0	24.3	0.0	60.6	100.0
Middle	0.0	11.7	0.0	1.6	7.3	0.0	41.6	0.0	37.7	100.0
Higher	0.0	2.7	8.0	1.9	6.0	1.7	25.0	8.3	46.3	100.0
Highest	0.0	3.7	8.0	7.3	5.5	0.0	24.6	3.9	47.0	100.0


¹Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Current Method Mix Among All and Married Contraceptive Users (Age 15-49)



Unmet need for family planning

Percentage of women age 15 to 49 with unmet need for family planning, by marital status and background characteristics

Background characteristics	All women			Married women ¹		
	Number	1,309		674		
	For spacing	For limiting	Total	For spacing	For limiting	Total
Total	9.8	4.6	14.4	12.1	8.3	20.4
Age group						
15–19	11.4	0.2	11.6	(38.7)	(0.0)	(38.7)
20–24	11.9	2.3	14.2	14.7	6.2	20.9
25–29	13.8	0.8	14.6	23.8	1.6	25.3
30–34	8.6	4.9	13.5	10.5	6.4	16.9
35–39	8.6	10.6	19.3	9.4	12.8	22.2
40–44	6.7	13.8	20.5	7.3	15.8	23.1
45–49	0.5	6.6	7.1	0.0	8.5	8.5
Marital status						
Married ¹	12.1	8.3	20.4	na	na	na
Unmarried, sexually active	31.2	2.6	33.8	na	na	na
Parity						
0–1	9.8	0.2	10.0	16.1	0.0	16.1
2–3	13.8	6.3	20.1	14.5	6.3	20.8
4 or more	6.9	13.4	20.3	7.9	15.0	22.9
Residence						
Urban	9.0	4.4	13.4	11.2	8.6	19.8
Rural	11.9	5.0	16.9	14.3	7.6	21.9
Education						
Never attended	(0.0)	(9.5)	(9.5)	(0.0)	(17.1)	(17.1)
Primary	3.6	9.9	13.4	4.9	14.4	19.3
Secondary	10.6	5.0	15.5	11.6	9.5	21.1
Higher	10.1	2.3	12.4	15.4	4.2	19.6
Wealth quintile						
Lowest	9.6	6.3	15.8	14.1	10.5	24.5
Lower	12.0	4.5	16.5	16.9	9.8	26.7
Middle	10.2	2.2	12.5	12.8	4.0	16.7
Higher	9.9	4.2	14.1	9.0	7.3	16.4
Highest	7.5	5.9	13.4	9.6	10.6	20.2

¹Married includes married women and those who are living with a man

na = not applicable

Need and demand for family planning

Percentage of women age 15 to 49 with met need, unmet need, and total demand for family planning, and percentage of demand for family planning satisfied with modern methods, by marital status and background characteristics

Background characteristics Number	All women 1,309					Married women ² 674				
	Any Method	Modern method ¹	Unmet need	Total demand	Demand satisfied by modern method	Any Method	Modern method ¹	Unmet need	Total demand	Demand satisfied by modern method
Total	25.1	15.6	14.4	39.5	39.6	33.5	17.5	20.4	53.9	32.6
Age group										
15–19	6.9	6.2	11.6	18.5	33.5	(17.4)	(17.4)	(38.7)	(56.1)	(31.0)
20–24	24.9	21.3	14.2	39.1	54.6	26.6	20.0	20.9	47.5	42.2
25–29	26.7	17.6	14.6	41.4	42.5	26.3	14.5	25.3	51.6	28.2
30–34	33.2	16.6	13.5	46.7	35.6	38.8	17.5	16.9	55.7	31.3
35–39	38.8	20.4	19.3	58.1	35.2	41.7	20.6	22.2	63.9	32.2
40–44	24.2	15.1	20.5	44.7	33.9	28.5	17.8	23.1	51.6	34.5
45–49	25.5	11.1	7.1	32.6	33.9	34.7	14.4	8.5	43.2	33.3
Marital status										
Married ²	33.5	17.5	20.4	53.9	32.6	na	na	na	na	na
Unmarried, sexually active	36.7	31.2	33.8	70.5	44.2	na	na	na	na	na
Parity										
0–1	15.4	12.3	10.0	25.3	48.5	13.9	9.0	16.1	30.0	30.2
2–3	29.3	17.6	20.1	49.4	35.5	29.7	16.8	20.8	50.5	33.3
4 or more	44.7	22.2	20.3	65.0	34.1	48.8	23.5	22.9	71.7	32.8
Residence										
Urban	26.4	16.2	13.4	39.8	40.6	35.1	17.3	19.8	54.9	31.6
Rural	21.7	14.2	16.9	38.6	36.7	29.6	18.1	21.9	51.4	35.1
Education										
Never attended	(7.0)	(0.0)	(9.5)	(16.5)	(0.0)	(0.0)	(0.0)	(17.1)	(17.1)	(0.0)
Primary	25.1	11.9	13.4	38.6	30.7	33.9	16.4	19.3	53.2	30.8
Secondary	22.6	13.9	15.5	38.1	36.6	33.2	17.4	21.1	54.3	32.0
Higher	30.8	20.5	12.4	43.2	47.5	34.6	18.5	19.6	54.2	34.2
Wealth quintile										
Lowest	19.2	13.5	15.8	35.1	38.5	28.2	18.8	24.5	52.7	35.6
Lower	20.2	11.2	16.5	36.7	30.5	28.6	11.3	26.7	55.3	20.4
Middle	26.8	18.6	12.5	39.3	47.2	32.6	20.3	16.7	49.3	41.1
Higher	31.3	18.1	14.1	45.4	39.8	37.6	17.6	16.4	53.9	32.7
Highest	26.5	16.3	13.4	39.9	40.9	37.4	19.2	20.2	57.7	33.3

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

Pregnancy intentions

Unintended births among women age 15 to 49 with one or more births in the previous 5 years or who are currently pregnant: Percent distribution reporting last or current pregnancy was wanted then, wanted later or not wanted at all, by marital status and background characteristics

Background characteristics	All women with 1 or more births				Married women with 1 or more births			
	469				433			
Number	Wanted then	Wanted later	Wanted no more	Total	Wanted then	Wanted later	Wanted no more	Total
Total	71.5	16.7	11.8	100.0	73.6	16.3	10.0	100.0
Age group								
15–19	(51.7)	(35.3)	(13.0)	(100.0)	(60.2)	(39.8)	(0.0)	(100.0)
20–24	65.5	24.9	9.5	100.0	70.7	24.4	4.8	100.0
25–29	73.3	20.0	6.8	100.0	77.7	18.7	3.6	100.0
30–34	71.8	15.6	12.6	100.0	71.7	15.8	12.6	100.0
35–39	73.1	10.6	16.3	100.0	73.5	9.8	16.7	100.0
40–44	78.9	9.4	11.8	100.0	80.4	10.9	8.7	100.0
45–49	(45.3)	(0.0)	(54.7)	(100.0)	(45.6)	(0.0)	(54.7)	(100.0)
Marital status								
Married ¹	73.6	16.3	10.0	100.0	na	na	na	na
Unmarried, sexually active	(46.7)	(30.8)	(22.5)	(100.0)	na	na	na	na
Parity								
0–1	72.2	20.3	7.5	100.0	78.0	20.6	1.4	100.0
2–3	80.5	15.0	4.5	100.0	82.5	13.6	3.9	100.0
4 or more	61.4	16.0	22.6	100.0	61.1	16.6	22.3	100.0
Residence								
Urban	76.1	12.6	11.2	100.0	77.4	12.2	10.5	100.0
Rural	61.1	25.8	13.1	100.0	64.7	26.4	8.9	100.0
Education								
Never attended	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)	(0.0)	(0.0)	(100.0)
Primary	74.8	10.3	14.8	100.0	76.3	11.8	11.9	100.0
Secondary	67.7	18.1	14.2	100.0	70.1	17.6	12.4	100.0
Higher	77.4	15.9	6.6	100.0	79.2	15.5	5.3	100.0
Wealth quintile								
Lowest	68.6	21.3	10.1	100.0	72.8	21.0	6.2	100.0
Lower	73.5	16.3	10.2	100.0	74.8	15.6	9.7	100.0
Middle	58.3	20.4	21.3	100.0	60.1	22.1	17.8	100.0
Higher	77.9	15.5	6.6	100.0	79.5	13.5	6.9	100.0
Highest	79.3	10.9	9.7	100.0	80.6	11.2	8.2	100.0

¹Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Contraceptive choice

Percent distribution of women age 15 to 49 who are currently using modern contraception¹ reporting on who decided on the contraceptive method, by background characteristics

Background characteristics	Self	Jointly with partner or provider	Provider alone/ partner alone/other	Total
Number	82	198	46	326
Total	25.2	60.6	14.2	100.0
Age group				
15–19	(37.0)	(54.6)	(8.4)	(100.0)
20–24	32.4	53.5	14.1	100.0
25–29	24.7	44.9	30.3	100.0
30–34	28.9	64.7	6.5	100.0
35–39	22.3	69.2	8.5	100.0
40–44	8.9	76.7	14.4	100.0
45–49	(19.7)	(66.7)	(13.1)	(100.0)
Marital status				
Married ²	18.3	66.7	15.0	100.0
Unmarried, sexually active	44.4	43.6	12.0	100.0
Parity				
0–1	33.8	51.9	14.3	100.0
2–3	19.6	65.0	15.3	100.0
4 or more	21.3	65.2	13.5	100.0
Residence				
Urban	26.4	61.1	12.5	100.0
Rural	21.1	59.2	19.7	100.0
Education				
Never attended	(100.0)	(0.0)	(0.0)	(100.0)
Primary	24.1	49.5	26.3	100.0
Secondary	25.0	63.6	11.4	100.0
Higher	25.4	58.9	15.8	100.0
Wealth quintile				
Lowest	27.9	51.8	20.3	100.0
Lower	23.9	61.1	15.1	100.0
Middle	27.1	55.4	17.5	100.0
Higher	23.1	69.2	7.7	100.0
Highest	25.3	59.9	14.8	100.0

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Payment for family planning

Percentage of women age 15 to 49 currently using a modern contraceptive method¹ who paid fees for services, by marital status and background characteristics

Background characteristics	All modern contraceptive users ¹	Married ² modern contraceptive users ¹
Number	329	226
Total	29.0	23.7
Age group		
15–19	40.2	(65.5)
20–24	42.3	(32.0)
25–29	26.3	18.8
30–34	30.5	24.2
35–39	28.7	30.7
40–44	21.4	21.4
45–49	4.9	(5.2)
Marital status		
Married ²	23.7	na
Unmarried, sexually active	47.9	na
Parity		
0–1	37.1	(26.9)
2–3	25.4	22.2
4 or more	24.4	23.9
Residence		
Urban	27.9	21.3
Rural	32.7	30.8
Education		
Never attended	(0.0)	(0.0)
Primary	36.1	(35.4)
Secondary	28.4	20.3
Higher	28.5	26.1
Wealth quintile		
Lowest	32.0	24.0
Lower	31.4	26.7
Middle	28.3	27.9
Higher	29.2	21.3
Highest	26.2	21.0

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Method information Index Components

Percentage of current modern¹ contraceptive users who reported whether provider informed them about other methods, side effects and, if informed of side effects, what to do, by marital status and background characteristics²

Background characteristics	Informed about other methods		Informed about side effects		Told what to do if experienced side effects	
	All users	Married users ³	All users	Married users ³	All users	Married users ³
Number	329	226	208	122	52	35
Total	36.3	39.4	25.1	28.6	68.6	70.6
Age group						
15–19	(5.7)	(0.0)	(0.0)	(0.0)	na	na
20–24	25.5	19.2	18.6	(9.2)	(33.9)	(0.0)
25–29	40.2	42.1	19.6	(23.1)	(100)	(100.0)
30–34	28.6	29.9	32.1	37.0	(89.7)	(88.9)
35–39	43.9	42.6	35.3	28.0	(62.4)	(59.0)
40–44	52.1	52.1	(24.0)	(24.0)	(100.0)	(100.0)
45–49	(52.7)	(56.2)	(52.0)	(57.5)	(26.0)	(26.0)
Marital status						
Married ³	39.4	na	28.6	na	(70.6)	na
Unmarried, sexually active	20.9	na	15.3	na	(35.2)	na
Parity						
0–1	23.3	(7.3)	13.9	(0.0)	(63.2)	na
2–3	42.5	41.7	26.2	24.2	(46.4)	(49.3)
4 or more	43.5	43.9	38.0	37.5	79.8	(80.0)
Residence						
Urban	33.6	36.6	27.5	33.9	73.70	(76.7)
Rural	45.0	47.5	17.5	15.8	(43.7)	(39.3)
Education						
Never attended	(0.0)	na	na	na	na	na
Primary	46.3	(48.6)	(10.2)	(11.6)	(100.0)	(100.0)
Secondary	25.6	30.0	22.0	24.5	(73.8)	(79.2)
Higher	50.3	52.5	31.7	40.1	(62.0)	(59.7)
Wealth quintile						
Lowest	34.1	40.7	4.4	(2.3)	(65.2)	(0.0)
Lower	30.3	35.4	12.8	(16.1)	(40.6)	(65.4)
Middle	29.3	29.4	28.7	(29.2)	(88.8)	(100.0)
Higher	38.7	37.5	22.0	26.4	(43.0)	(56.2)
Highest	45.4	50.9	45.4	54.0	(74.1)	(65.1)

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²"Informed about other methods" asked of all current contraceptive users, "informed about side effects" asked of current modern contraceptive users, and "told what to do if experienced side effects" asked of current modern contraceptive users who indicated they were told about side effects.

³Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Perceived quality of care for family planning

Percentage of current modern¹ contraceptive users who would return, would refer a friend/relative, or would return and refer friend/relative to provider, by background characteristics

Background characteristics	Percent who would return to provider	Percent who would refer relative/friend	Percent who would return and refer friend/relative
Number	208	208	208
Total	55.7	54.0	43.5
Age group			
15–19	(43.0)	(17.8)	(17.8)
20–24	41.4	40.2	29.4
25–29	65.2	64.4	53.6
30–34	57.5	60.8	43.3
35–39	52.3	61.8	49.9
40–44	(67.8)	(52.3)	(47.0)
45–49	(80.9)	(77.5)	(73.0)
Marital status			
Married ²	63.2	64.3	53.2
Unmarried, sexually active	44.7	38.7	25.4
Parity			
0–1	45.6	37.1	29.0
2–3	57.4	68.2	51.7
4 or more	67.1	66.4	56.5
Residence			
Urban	55.3	52.2	41.2
Rural	57.0	59.7	50.6
Education			
Never attended	na	na	na
Primary	(73.3)	(61.0)	(55.1)
Secondary	56.4	52.5	43.2
Higher	52.1	55.1	42.2
Wealth quintile			
Lowest	53.7	46.4	42.9
Lower	47.8	52.9	43.0
Middle	54.6	36.9	32.9
Higher	50.7	58.2	42.1
Highest	68.8	72.7	56.9

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

Knowledge of family planning

Percentage of women age 15 to 49 who reported receiving family planning information from a health worker in the past 12 months, by marital status and background characteristics

Background characteristics	Visited by health worker who discussed family planning		Visited facility and provider discussed family planning		Either visited by health worker or visited facility and discussed family planning	
	All women	Married women ¹	All women	Married women ¹	All women	Married women ¹
Number	1,309	674	1,309	674	1,309	674
Total	11.4	16.4	17.1	25.8	21.9	32.1
Age group						
15–19	3.1	(0.0)	2.3	(18.2)	5.1	(18.2)
20–24	9.0	20.2	13.2	22.8	16.9	29.8
25–29	14.3	21.7	24.0	37.6	28.5	44.2
30–34	13.5	14.6	25.8	28.7	29.4	33.4
35–39	14.3	13.7	22.6	22.9	29.9	30.3
40–44	13.0	14.6	20.8	22.1	24.2	26.1
45–49	17.7	18.3	12.2	13.9	24.6	25.2
Marital status						
Married ¹	16.4	na	25.8	na	32.1	na
Unmarried, sexually active	8.4	na	15.1	na	17.8	na
Parity						
0–1	7.2	12.4	10.9	22.8	13.7	25.6
2–3	18.8	19.7	31.6	33.8	38.5	40.0
4 or more	15.3	16.4	20.6	21.7	28.4	30.2
Residence						
Urban	11.6	16.2	17.0	25.9	21.3	31.1
Rural	10.8	16.8	17.4	25.7	23.4	34.7
Education						
Never attended	(0.0)	(0.0)	(7.0)	(12.6)	(7.0)	(12.6)
Primary	4.6	6.5	14.6	18.1	17.2	22.2
Secondary	9.8	14.1	15.8	25.2	19.6	30.1
Higher	16.5	23.8	20.7	29.5	28.2	39.2
Wealth quintile						
Lowest	7.5	11.3	10.1	16.4	14.0	22.9
Lower	14.0	19.3	14.4	19.5	20.3	27.9
Middle	8.9	13.3	19.6	26.6	22.9	31.9
Higher	13.7	21.0	22.1	32.5	27.7	39.7
Highest	11.7	15.3	17.6	29.3	22.6	33.9

¹Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

List of indicators for sampling errors, PMA2016/Anambra

Variable	Estimation	Base population
Currently using a modern method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using a traditional method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using any contraceptive method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using injectables	Proportion	All women ages 15–49 Married women ages 15–49
Currently using male condoms	Proportion	All women ages 15–49 Married women ages 15–49
Currently using implants	Proportion	All women ages 15–49 Married women ages 15–49
Chose method by self or jointly in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49
Paid fees for family planning services in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49
Informed by provider about other methods	Proportion	All women ages 15–49 Married women ages 15–49
Informed by provider about side effects	Proportion	All women ages 15–49 Married women ages 15–49
Satisfied with provider: Would return and refer friend/relative to provider	Proportion	All women ages 15–49 Married women ages 15–49
Visited by health worker who talked about family planning information in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49

Characteristics of service delivery points

Number and percentage of service delivery points (SDPs) by background characteristics

Background characteristics	Number	Percentage
Total	102	100.0
SDP type		
Hospital	9	8.8
Health center ¹	44	43.1
Pharmacy ²	48	47.1
Other	1	1.0
Sector		
Public	51	50.0
Private	51	50.0
Offers family planning		
No	38	37.3
Yes	64	62.7

¹Includes Health Post, Health Clinic, and Maternity Center

²Includes Chemist and Patent Medicine Vendors

Charging fees for family planning

Percentage of service delivery points (SDPs) that charge fees for family planning services, by sector and background characteristics, among SDPs offering family planning

Background characteristic	Sector	
	Public	Private
Number of SDPs	32	32
Total	50.0	53.1
SDP type		
Hospital	(0.0)	(100.0)
Health center ¹	55.2	na
Pharmacy ²	na	51.6
Other	na	na

¹Includes Health Post, Health Clinic, and Maternity Center

²Includes Chemist and Patent Medicine Vendors

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Contraceptive choice: Availability of at least 3 or 5 modern contraceptive methods

Percentage of public and private service delivery points (SDPs) offering at least 3 or 5 modern contraceptive methods¹, by sector and background characteristics, among all SDPs

Background characteristic	3 or more methods		5 or more methods	
	Public	Private	Public	Private
Number of SPDs	51	51	51	51
Total	29.4	9.8	13.7	2.0
SDP type				
Hospital	(12.5)	(0.0)	(0.0)	(0.0)
Health Center ²	33.3	(0.0)	16.7	(0.0)
Pharmacy ³	na	10.4	na	2.1
Other	(0.0)	na	(0.0)	na

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Includes Health Post, Health Clinic, and Maternity Center

³Includes Chemist and Patent Medicine Vendors

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Contraceptive Choice: Availability of modern contraception, by method

Percentage of service delivery points (SDPs) offering different types of contraceptive methods¹, by sector and method, among all SDPs

Public SDPs							
Background characteristic	Contraceptive method						
	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception
Number of SDPs = 51							
Total	49.0	52.9	13.7	7.8	47.1	0.0	5.9
SDP type							
Hospital	(12.5)	(12.5)	(12.5)	(0.0)	(12.5)	(0.0)	(0.0)
Health center ²	57.1	61.9	14.3	9.5	54.8	0.0	7.1
Pharmacy ³	na	na	na	na	na	na	na
Other	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)

Private SDPs							
Background characteristic	Contraceptive method						
	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception
Number of SDPs = 51							
Total	27.5	3.9	2.0	0.0	56.9	2.0	9.8
SDP type							
Hospital	(100.0)	(0.0)	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
Health center ²	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)
Pharmacy ³	27.1	4.2	0.0	0.0	60.4	0.0	8.3
Other	na	na	na	na	na	na	na

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Includes Health Post, Health Clinic, and Maternity Center

³Includes Chemist and Patent Medicine Vendors

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Contraceptive stock-outs, by method

Percentage of service delivery points (SDPs) stocked out of specific modern contraceptives¹ at time of survey or stocked out in the last 3 months, by sector and background characteristics, among SDPs that offer each method

Public SDPs						
Background characteristic	Contraceptive method					
	Pill	Injectables	IUD	Implants	Male condom	Emergency contraception
Number of SDPs	25	27	7	4	24	3
Total	20.0	22.2	(14.3)	(25.0)	(41.7)	(0.0)
SDP type						
Hospital	(0.0)	(0.0)	(0.0)	na	(0.0)	na
Health center ¹	(20.8)	23.1	(16.7)	(25.0)	(43.5)	(0.0)
Pharmacy ²	na	na	na	na	na	na
Other	na	na	na	na	na	na

Private SDPs						
Background characteristic	Contraceptive method					
	Pill	Injectables	IUD	Implants	Male condom	Emergency contraception
Number of SDPs	14	2	1	0	29	5
Total	(21.4)	(50.0)	(100.0)	na	20.7	(40.0)
SDP type						
Hospital	(100.0)	na	(100.0)	na	na	(100.0)
Health center ¹	na	na	na	na	na	na
Pharmacy ²	(15.4)	(50.0)	na	na	20.7	(25.0)
Other	na	na	na	na	na	na

¹Includes Health Post, Health Clinic, and Maternity Center

²Includes Chemist and Patent Medicine Vendors

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Number of family planning visits, by method

Total number of visits and new clients contraceptive services, by method type and background characteristics, among advanced¹ service delivery points (SDPs) that offer each method

	Contraceptive Methods													
	Sterilization		Injectable		IUD		Implants		Male condom		Pills		Emergency Contraception	
	Total women	Total men	Total	New	Total	New	Total	New	Total	New	Total	New	Total	New
Number of SDPs	1	0	27		8		4		24		26		4	
Total	na	na	191	56	72	38	51	28	148	66	118	54	58	36
Background characteristic														
SDP type														
Hospital	na	na	4	4	0	0	0	0	18	18	8	8	0	0
Health center ¹	na	na	187	52	72	38	51	28	130	48	110	46	58	36
Sector														
Public	na	na	191	56	72	38	51	28	148	66	118	54	58	36
Private	na	na	0	0	0	0	0	0	0	0	0	0	0	0

¹Includes Health Post, Health Clinic, and Maternity Center