Background characteristics of respondents

Percent distribution of background characteristics of women age 15 to 49

Background characteristics	Urban	Rural	Percentage of sample	Weighted number	Unweighted number
Age group					
15–19	22.3	77.7	100.0	18.5	303
20–24	18.1	81.9	100.0	21.3	348
25–29	20.7	79.3	100.0	20.3	332
30–34	20.4	79.6	100.0	16.4	269
35–39	20.7	79.3	100.0	9.5	156
40–44	19.1	80.9	100.0	9.2	150
45–49	25.9	74.1	100.0	4.8	79
Marital status					
Married	20.3	79.7	100.0	68.2	1,117
Living together	(15.7)	(84.3)	(100.0)	0.8	14
Divorced	22.8	77.2	100.0	2.1	34
Widowed	18.4	81.6	100.0	3.4	55
Never married	21.3	78.7	100.0	25.4	416
Parity					
None	21.4	78.6	100.0	28.6	469
1-2	19.6	80.4	100.0	28.1	460
3-4	18.6	81.4	100.0	22.1	362
5 or more	22.5	77.5	100.0	21.1	346
Education					
Never attended	10.7	89.3	100.0	25.4	416
Primary	23.8	76.2	100.0	24.8	407
Secondary	18.8	81.2	100.0	39.4	645
Higher	42.7	57.3	100.0	10.4	171
Wealth quintile					
Lowest	0.4	99.6	100.0	20.8	341
Lower	4.9	95.1	100.0	20.9	342
Middle	15.6	84.4	100.0	19.0	311
Higher	37.7	62.3	100.0	19.7	323
Highest	45.8	54.2	100.0	19.6	321
Total	**20.5**	**79.5**	**100.0**	**100.0**	**1,638**

PMA2020 Performance Monitoring and Accountability 2020

© 2017 Performance Monitoring and Accountability 2020 (PMA2020)

Current use of contraception by background characteristics

Percentage of women ages 15 to 49 currently using a contraceptive method, by type and background characteristics

Background characteristic	Any method		Modern method[1]		Traditional method	
	All women	Married women	All women	Married women	All women	Married women
N	1,638	1,130	1,638	1,130	1,638	1,130
Total	**18.9**	**21.5**	**16.6**	**18.9**	**2.4**	**2.6**
Age group						
15–19	6.1	11.0	5.6	11.0	0.5	0.0
20–24	15.3	15.1	10.8	9.6	4.5	5.5
25–29	17.5	16.7	14.1	12.9	3.3	3.9
30–34	28.5	27.1	25.3	25.2	3.2	1.9
35–39	31.4	31.1	31.0	30.6	0.4	0.5
40–44	27.9	29.3	27.7	29.0	0.2	0.3
45–49	16.0	19.2	15.1	18.7	0.9	0.5
Marital status						
Married[2]	21.5	na	18.9	na	2.6	na
Unmarried, sexually active	43.8	na	38.5	na	5.2	na
Parity						
0–1	12.4	12.2	9.5	7.9	2.9	4.2
2–3	18.8	19.3	16.8	17.0	2.1	2.3
4 or more	28.3	29.0	26.5	27.2	1.9	1.8
Residence						
Urban	15.6	20.6	13.2	17.1	2.4	3.5
Rural	19.8	21.7	17.4	19.4	2.4	2.4
Education						
Never attended	13.8	15.0	12.0	13.2	1.9	1.8
Primary	20.3	22.8	17.5	19.7	2.7	3.1
Secondary	20.6	25.8	18.5	23.2	2.2	2.6
Higher	21.7	27.2	18.5	23.3	3.2	3.9
Wealth quintile						
Lowest	18.2	18.2	15.7	16.9	2.5	1.3
Lower	19.8	25.0	18.4	23.2	1.5	1.8
Middle	19.0	18.0	16.0	14.5	3.0	3.5
Higher	16.6	20.4	13.4	15.9	3.2	4.5
Highest	21.2	26.5	19.4	24.6	1.7	1.8

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

Contraceptive method mix by background characteristics

Percent distribution of current contraceptive users ages 15 to 49, by method type, marital status and background characteristics

Background characteristics	Steriliz-ation	Implants	IUD	Inject-ables	Pill	Emergency contraception	Condom	Other modern	Other traditional	Total
All users Number = 310										
Total	**2.4**	**25.3**	**3.9**	**30.4**	**9.2**	**1.2**	**14.7**	**1.4**	**11.6**	**100.0**
Age group										
15–19	(0.0)	(0.0)	(0.0)	(13.0)	(29.7)	(9.6)	(39.9)	(0.0)	(7.9)	(100.0)
20–24	0.0	8.8	0.0	18.5	19.2	3.4	19.3	3.5	27.2	100.0
25–29	2.4	35.4	0.0	23.0	9.4	0.0	10.8	1.9	17.1	100.0
30–34	0.0	23.0	5.5	36.4	3.2	0.0	20.6	0.0	11.4	100.0
35–39	3.7	40.3	5.5	40.3	4.3	0.0	4.5	0.0	1.4	100.0
40–44	10.4	32.2	4.4	34.9	6.6	0.0	8.4	2.3	0.8	100.0
45–49	(0.0)	(17.1)	(26.6)	(50.8)	(0.0)	(0.0)	(0.0)	(2.8)	(2.7)	(100.0)
Marital status										
Married[1]	2.3	28.8	4.5	36.2	6.9	0.0	8.6	1.6	11.1	100.0
Unmarried, sexually active	0.0	11.2	0.0	11.8	21.2	0.0	43.8	0.0	11.9	100.0
Parity										
0–1	0.0	8.8	0.0	8.1	20.4	4.0	35.4	2.5	20.8	100.0
2–3	1.9	24.1	2.0	42.5	5.5	0.0	12.0	1.0	10.9	100.0
4 or more	4.2	35.9	7.3	37.9	4.2	0.0	3.2	0.9	6.3	100.0
Residence										
Urban	4.7	24.3	6.1	24.8	8.9	3.4	10.8	6.1	10.9	100.0
Rural	2.0	25.4	3.5	31.5	9.2	0.7	15.4	0.4	11.7	100.0
Education										
Never attended	0.0	20.0	2.0	59.0	5.4	0.0	0.0	0.6	12.9	100.0
Primary	7.0	29.7	2.0	34.3	4.7	0.0	7.9	0.9	13.5	100.0
Secondary	1.4	29.1	5.3	17.3	12.0	2.7	20.9	1.6	9.8	100.0
Higher	0.0	9.5	6.0	24.3	15.1	0.0	30.1	3.0	11.9	100.0
Wealth quintile										
Lowest	5.3	14.5	3.3	43.9	9.2	0.0	10.0	0.0	13.8	100.0
Lower	2.7	30.7	4.4	34.3	9.9	2.7	6.7	1.1	7.4	100.0
Middle	0.0	21.3	0.0	24.4	8.8	0.0	29.8	4.4	11.3	100.0
Higher	4.6	40.0	1.8	18.9	7.3	3.3	5.0	0.0	19.1	100.0
Highest	0.0	21.5	9.0	28.4	10.3	0.0	21.1	1.4	8.2	100.0

[1]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

| Background characteristics	Steriliz-ation	Implants	IUD	Inject-ables	Pill	Emergency contraception	Condom	Other modern	Other traditional	Total
Married users										
Number = 243										
Total	**2.3**	**28.8**	**4.5**	**36.2**	**6.9**	**0.0**	**8.6**	**1.6**	**11.1**	**100.0**
Age group										
15–19	(0.0)	(0.0)	(0.0)	(33.3)	(25.1)	(0.0)	(41.6)	(0.0)	(0.0)	(100.0)
20–24	0.0	13.2	0.0	23.6	13.7	0.0	10.8	5.3	33.5	100.0
25–29	3.1	33.3	0.0	27.4	8.1	0.0	5.0	2.5	20.6	100.0
30–34	0.0	28.9	6.9	44.4	2.9	0.0	9.8	0.0	7.0	100.0
35–39	4.3	39.9	3.6	40.4	4.9	0.0	5.2	0.0	1.6	100.0
40–44	6.3	34.5	4.7	35.6	6.5	0.0	9.1	2.5	0.9	100.0
45–49	(0.0)	(17.6)	(27.4)	(52.2)	(0.0)	(0.0)	(0.0)	(0.0)	(2.8)	(100.0)
Marital status										
Married[1]	na	na	na	na	na	na	na	na	na	na
Unmarried, sexually active	na	na	na	na	na	na	na	na	na	na
Parity										
0–1	0.0	13.3	0.0	11.8	18.8	0.0	21.2	6.2	28.7	100.0
2–3	2.0	21.7	2.2	43.5	5.6	0.0	12.3	1.1	11.7	100.0
4 or more	3.1	36.4	6.9	38.8	4.4	0.0	3.4	0.7	6.2	100.0
Residence										
Urban	5.2	19.5	6.8	27.6	9.9	0.0	12.0	6.8	12.1	100.0
Rural	1.6	31.0	4.0	38.2	6.2	0.0	7.8	0.4	10.8	100.0
Education										
Never attended	0.0	20.7	2.1	59.7	5.5	0.0	0.0	0.0	12.0	100.0
Primary	5.1	30.4	2.2	34.9	4.9	0.0	7.8	1.0	13.6	100.0
Secondary	2.1	36.9	8.1	24.5	5.8	0.0	11.5	2.4	8.7	100.0
Higher	(0.0)	(13.9)	(4.3)	(28.0)	(19.4)	(0.0)	(20.0)	(4.4)	(9.9)	(100.0)
Wealth quintile										
Lowest	3.1	19.7	4.5	53.5	7.8	0.0	4.2	0.0	7.2	100.0
Lower	3.6	37.5	3.6	38.6	2.4	0.0	5.6	1.5	7.1	100.0
Middle	0.0	25.9	0.0	34.6	7.5	0.0	12.4	5.3	14.4	100.0
Higher	5.5	36.9	2.2	22.7	7.2	0.0	3.5	0.0	22.1	100.0
Highest	0.0	24.2	10.2	32.1	9.4	0.0	15.6	1.6	6.9	100.0

[1]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Current Method Mix Among **All and Married** Contraceptive Users (Age 15-49)

Number of users:
310

Number of users:
243

Percentage

All

Value	Method
11.6	
1.4	
1.2	
14.7	
9.2	
30.4	
25.3	
3.9	
2.4	

Married

Value	Method
11.1	
1.6	
8.6	
6.9	
36.2	
28.8	
4.5	
2.3	

Legend:
- Other traditional
- Other modern
- Emergency contraception
- Male condoms
- Pills
- Injectables
- Implants
- IUD
- Female Sterilization

PMA2020 | Performance Monitoring and Accountability 2020

Unmet need for family planning

Percentage of women ages 15 to 49 with unmet need for family planning (for spacing or limiting), by marital status and background characteristics

Background characteristics	All women			Married women		
Number	**1,638**			**1,130**		
	For spacing	**For limiting**	**Total**	**For spacing**	**For limiting**	**Total**
Total	**11.9**	**6.0**	**18.0**	**13.9**	**8.1**	**22.0**
Age group						
15–19	8.8	1.9	10.7	14.3	1.0	15.3
20–24	14.8	4.4	19.2	16.5	6.4	22.9
25–29	14.6	3.6	18.2	16.1	4.5	20.5
30–34	16.4	8.7	25.1	18.9	10.4	29.3
35–39	11.1	11.6	22.7	11.1	13.3	24.4
40–44	4.0	9.8	13.8	4.5	10.7	15.2
45–49	1.8	11.7	13.4	2.2	12.7	14.9
Marital status						
Married[1]	13.9	8.1	22.0	na	na	na
Unmarried, sexually active	38.2	4.2	42.4	na	na	na
Parity						
0–1	8.9	0.6	9.5	10.4	0.0	10.4
2–3	17.5	6.4	23.9	18.4	6.5	24.9
4 or more	12.0	13.1	25.1	12.6	14.2	26.8
Residence						
Urban	10.5	5.5	16.0	12.6	7.6	20.3
Rural	12.3	6.1	18.5	14.2	8.3	22.4
Education						
Never attended	13.7	10.4	24.1	14.5	11.3	25.8
Primary	11.6	5.9	17.5	13.4	7.3	20.7
Secondary	10.7	3.7	14.5	12.6	5.8	18.4
Higher	13.1	4.4	17.4	17.5	6.8	24.3
Wealth quintile						
Lowest	12.8	6.6	19.4	14.7	9.0	23.7
Lower	13.4	4.8	18.3	15.6	7.1	22.7
Middle	10.0	8.0	18.0	12.7	10.7	23.4
Higher	11.9	5.4	17.2	14.0	7.9	21.9
Highest	11.4	5.4	16.8	12.5	5.6	18.1

[1]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

Need and demand for family planning

Percentage women ages 15 to 49 with met need, unmet need, and total demand for family planning, and percentage of demand for family planning satisfied with modern methods, by marital status and background characteristics

Background characteristics	All women					Married women				
Number	**1,638**					**1,130**				
	Modern method[1]	Any method	Unmet need	Total demand	Demand satisfied by modern method	Modern method[2]	Any method	Unmet need	Total demand	Demand satisfied by modern method
Total	**16.6**	**18.9**	**18.0**	**36.9**	**44.9**	**18.9**	**21.5**	**22.0**	**43.5**	**43.5**
Age group										
15–19	5.6	6.1	10.7	16.8	33.4	11.0	11.0	15.3	26.3	(41.8)
20–24	10.8	15.3	19.2	34.5	31.4	9.6	15.1	22.9	38.0	25.2
25–29	14.1	17.5	18.2	35.7	39.6	12.9	16.7	20.5	37.3	34.5
30–34	25.3	28.5	25.1	53.6	47.1	25.2	27.1	29.3	56.4	44.6
35–39	31.0	31.4	22.7	54.1	57.2	30.6	31.1	24.4	55.6	55.1
40–44	27.7	27.9	13.8	41.8	66.3	29.0	29.3	15.2	44.5	65.2
45–49	15.1	16.0	13.4	29.4	51.4	18.7	19.2	14.9	34.1	54.9
Marital status										
Married[2]	18.9	21.5	22.0	43.5	43.5	na	na	na	na	na
Unmarried, sexually active	38.5	43.8	42.4	86.2	44.7	na	na	na	na	na
Parity										
0–1	9.5	12.4	9.5	21.9	43.3	7.9	12.2	10.4	22.5	35.2
2–3	16.8	18.8	23.9	42.7	39.3	17.0	19.3	24.9	44.2	38.5
4 or more	26.5	28.3	25.1	53.5	49.5	27.2	29.0	26.8	55.9	48.7
Residence										
Urban	13.2	15.6	16.0	31.6	41.9	17.1	20.6	20.3	40.8	41.8
Rural	17.4	19.8	18.5	38.3	45.6	19.4	21.7	22.4	44.1	43.9
Education										
Never attended	12.0	13.8	24.1	37.9	31.5	13.2	15.0	25.8	40.8	32.3
Primary	17.5	20.3	17.5	37.8	46.4	19.7	22.8	20.7	43.6	45.3
Secondary	18.5	20.6	14.5	35.1	52.6	23.2	25.8	18.4	44.1	52.6
Higher	18.5	21.7	17.4	39.2	47.2	23.3	27.2	24.3	51.5	45.3
Wealth quintile										
Lowest	15.7	18.2	19.4	37.5	41.7	16.9	18.2	23.7	42.0	40.3
Lower	18.4	19.8	18.3	38.1	48.2	23.2	25.0	22.7	47.7	48.6
Middle	16.0	19.0	18.0	37.0	43.3	14.5	18.0	23.4	41.4	34.9
Higher	13.4	16.6	17.2	33.8	39.7	15.9	20.4	21.9	42.4	37.6
Highest	19.4	21.2	16.8	38.0	51.1	24.6	26.5	18.1	44.6	55.3

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

Pregnancy intentions

Unintended births among women ages 15 to 49 with one or more births in the previous 5 years or who are currently pregnant: Percent distribution reporting last or current pregnancy was wanted then, wanted later or not wanted at all, by marital status and background characteristics

Background characteristics	All women with 1 or more births				Married women with 1 or more births			
Number	887				822			
	Wanted then	Wanted later	Not at all	Total	Wanted then	Wanted later	Not at all	Total
Total	**70.0**	**22.0**	**8.0**	**100.0**	**71.9**	**21.4**	**6.7**	**100.0**
Age group								
15–19	75.4	16.1	8.5	100.0	85.2	14.8	0.0	100.0
20–24	70.5	21.9	7.6	100.0	73.2	20.8	6.0	100.0
25–29	71.1	25.7	3.2	100.0	73.6	24.6	1.8	100.0
30–34	69.5	24.9	5.6	100.0	69.4	25.4	5.3	100.0
35–39	69.9	21.0	9.1	100.0	70.7	19.7	9.6	100.0
40–44	62.3	11.2	26.6	100.0	61.2	11.5	27.3	100.0
45–49	62.3	0.0	37.7	100.0	56.5	0.0	43.5	100.0
Marital status								
Married[1]	71.9	21.4	6.7	100.0	na	na	na	na
Unmarried, sexually active	27.6	28.9	43.5	100.0	na	na	na	na
Parity								
0–1	79.0	15.7	5.2	100.0	86.1	13.9	0.0	100.0
2–3	72.2	22.9	4.9	100.0	73.3	22.0	4.7	100.0
4 or more	61.0	25.9	13.1	100.0	60.7	26.0	13.4	100.0
Residence								
Urban	67.7	25.6	6.7	100.0	68.7	25.4	5.9	100.0
Rural	70.6	21.1	8.3	100.0	72.7	20.4	6.9	100.0
Education								
Never attended	72.2	20.7	7.1	100.0	73.0	20.2	6.8	100.0
Primary	63.3	27.7	9.0	100.0	64.0	27.2	8.8	100.0
Secondary	71.1	19.1	9.8	100.0	75.6	18.0	6.5	100.0
Higher	80.0	20.0	0.0	100.0	80.2	19.8	0.0	100.0
Wealth quintile								
Lowest	73.9	21.2	4.9	100.0	75.5	20.5	4.0	100.0
Lower	65.0	27.3	7.6	100.0	67.4	26.0	6.6	100.0
Middle	77.7	15.8	6.6	100.0	78.4	15.3	6.3	100.0
Higher	70.7	20.2	9.2	100.0	74.5	19.3	6.2	100.0
Highest	60.8	26.7	12.5	100.0	61.9	27.1	11.0	100.0

[1]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

Contraceptive choice

Percent distribution of women ages 15 to 49 who are currently using modern contraception[1] reporting on who decided on the contraceptive method, by background characteristics

Background characteristics	Self	Jointly with partner or provider	Provider alone/ partner alone/other	Total
Number	**81**	**174**	**52**	**307**
Total	**26.5**	**56.7**	**16.8**	**100.0**
Age group				
15–19	(23.6)	(47.5)	(28.9)	(100.0)
20–24	40.0	50.7	9.3	100.0
25–29	30.2	49.0	20.9	100.0
30–34	25.5	61.1	13.4	100.0
35–39	12.8	73.2	14.0	100.0
40–44	16.5	63.6	19.9	100.0
45–49	(46.3)	(19.7)	(34.0)	(100.0)
Marital status				
Married[2]	18.7	64.2	17.1	100.0
Unmarried, sexually active	46.5	42.3	11.2	100.0
Parity				
0–1	44.5	36.1	19.4	100.0
2–3	10.8	68.1	21.1	100.0
4 or more	23.4	63.7	12.9	100.0
Residence				
Urban	27.6	61.2	11.3	100.0
Rural	26.3	55.8	17.9	100.0
Education				
Never attended	16.9	68.2	14.9	100.0
Primary	28.2	57.3	14.6	100.0
Secondary	29.1	51.5	19.4	100.0
Higher	28.7	56.2	15.2	100.0
Wealth quintile				
Lowest	25.5	64.8	9.7	100.0
Lower	27.5	47.1	25.4	100.0
Middle	24.8	52.7	22.5	100.0
Higher	28.0	56.7	15.3	100.0
Highest	26.7	62.7	10.6	100.0

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Payment for family planning

Percentage of women ages 15 to 49 currently using a modern contraceptive method[1] who paid fees for services, by marital status and background characteristics

Background characteristics	All women who used in past 12 months	All married women who used in past 12 month
Number	**310**	**243**
Total	**44.5**	**46.1**
Age group		
15–19	(33.2)	(4.7)
20–24	38.3	40.1
25–29	42.1	39.7
30–34	51.3	53.5
35–39	48.5	54.4
40–44	46.5	49.3
45–49	(34.8)	(35.8)
Marital status		
Married[2]	46.1	na
Unmarried, sexually active	50.6	na
Parity		
0–1	38.0	29.8
2–3	50.7	52.4
4 or more	45.3	47.2
Residence		
Urban	36.0	38.6
Rural	46.2	47.9
Education		
Never attended	55.8	57.7
Primary	51.0	52.2
Secondary	42.3	38.7
Higher	20.4	(28.1)
Wealth quintile		
Lowest	56.5	68.5
Lower	52.0	51.5
Middle	40.6	43.6
Higher	32.8	30.8
Highest	38.7	37.8

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Method Information Index Components

Percentage of current modern[1] contraceptive users who reported whether provider informed them about other methods, side effects and, if informed of side effects, what to do, by marital status and background characteristics[2]

Background characteristics	Informed about other methods		Informed about side effects		Told what to do if experienced side effects	
	All users	Married users	All users	Married users	All users	Married users
Number	310	243	287	224	128	111
Total	**51.3**	**56.4**	**44.5**	**49.3**	**86.3**	**92.0**
Age group						
15–19	(19.6)	(39.3)	(11.6)	(29.5)	*(100.0)*	*(100.0)*
20–24	43.5	62.8	18.3	27.6	(100.0)	(100.0)
25–29	43.3	39.4	56.3	54.4	75.7	(77.9)
30–34	64.9	64.1	56.1	62.0	84.4	94.4
35–39	51.2	53.9	61.0	63.5	95.0	94.5
40–44	59.7	64.0	29.8	27.1	(84.8)	(100.0)
45–49	(56.3)	(55.0)	(60.9)	(60.9)	(86.0)	(86.0)
Marital status						
Married[3]	56.4	na	49.3	na	92.0	na
Unmarried, sexually active	40.0	na	41.2	na	(59.2)	na
Parity						
0–1	44.4	61.5	27.2	(30.9)	(66.6)	(79.1)
2–3	65.9	67.4	57.1	58.0	93.0	94.5
4 or more	48.0	49.4	48.0	49.1	88.7	92.3
Residence						
Urban	48.3	51.4	52.1	56.2	87.2	(86.6)
Rural	51.9	57.6	43.1	47.7	86.1	93.4
Education						
Never attended	57.0	56.9	64.4	64.8	95.3	97.3
Primary	43.3	44.8	34.7	33.7	80.7	86.4
Secondary	48.7	60.4	40.2	47.6	78.3	87.5
Higher	69.2	(75.2)	49.7	(65.8)	(100.0)	(100.0)
Wealth quintile						
Lowest	43.4	47.7	37.7	38.5	84.8	100.0
Lower	48.3	56.6	54.4	65.9	89.0	91.0
Middle	53.7	59.2	38.3	54.3	(81.7)	(84.3)
Higher	52.6	52.8	35.0	31.6	(98.9)	(98.5)
Highest	58.2	63.3	52.2	51.8	(81.3)	(90.6)

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2] Informed about other methods" asked of all current contraceptive users, "informed about side effects" asked of current modern contraceptive users, and "told what to do if experienced side effects" asked of current modern contraceptive users who indicated they were told about side effects.

[3]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

PMA2020 Performance Monitoring and Accountability 2020

Perceived quality of care for family planning

Percentage of current modern[1] contraceptive users who would return, would refer a friend/relative, or would return and refer friend/relative to provider, by background characteristics

Background characteristics	Percent who would return to provider	Percent who would refer relative/friend	Percent who would return and refer friend/relative
Number	**287**	**287**	**287**
Total	**77.1**	**79.2**	**69.6**
Age group			
15–19	(79.0)	(70.8)	(61.3)
20–24	67.4	81.1	63.6
25–29	81.8	86.4	77.9
30–34	85.6	82.2	82.2
35–39	77.1	77.0	64.8
40–44	64.7	73.5	58.3
45–49	(85.2)	(66.0)	(54.3)
Marital status			
Married[2]	79.8	79.0	70.1
Unmarried, sexually active	82.0	91.4	75.3
Parity			
0–1	70.8	82.4	68.2
2–3	85.1	78.8	76.1
4 or more	76.7	77.6	67.3
Residence			
Urban	74.3	77.7	63.4
Rural	77.6	79.5	70.8
Education			
Never attended	78.8	73.2	69.9
Primary	69.2	71.0	59.6
Secondary	83.7	88.2	78.8
Higher	67.8	74.4	58.2
Wealth quintile			
Lowest	77.1	75.1	69.8
Lower	90.4	79.4	75.4
Middle	84.7	86.9	80.8
Higher	74.5	84.5	69.8
Highest	59.8	73.0	55.0

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Knowledge of family planning

Percentage of women ages 15 to 49 who reported receiving family planning information from a health worker in the past 12 months, by marital status and background characteristics

Background characteristics	Visited by health worker who discussed family planning		Visited facility and provider discussed family planning		Either visited by health worker or visited facility and discussed family planning	
	All women	Married women[1]	All women	Married women[1]	All women	Married women[1]
Number	**1,638**	**1,130**	**1,638**	**1,130**	**1,638**	**1,130**
Total	**13.1**	**17.0**	**18.7**	**24.4**	**25.0**	**32.4**
Age group						
15–19	5.5	14.6	6.6	23.3	9.8	29.9
20–24	12.6	14.5	17.5	23.1	24.3	31.0
25–29	18.2	21.3	26.0	31.5	31.3	37.3
30–34	16.7	20.0	22.9	23.8	33.6	36.6
35–39	18.6	19.7	22.9	21.2	31.0	30.5
40–44	9.3	10.4	19.1	21.4	24.2	27.2
45–49	7.2	9.0	15.8	14.9	20.1	20.2
Marital status						
Married[1]	17.0	na	24.4	na	32.4	na
Unmarried, sexually active	7.3	na	4.7	na	8.1	na
Parity						
0–1	9.4	16.8	11.7	23.3	16.5	31.5
2–3	18.3	19.3	27.7	29.6	34.9	36.9
4 or more	14.4	15.5	21.6	21.1	29.5	29.7
Residence						
Urban	14.3	18.0	26.4	35.8	31.8	42.0
Rural	12.8	16.8	16.7	21.5	23.2	30.0
Education						
Never attended	12.3	13.6	15.3	15.3	21.2	21.7
Primary	10.5	12.0	20.4	24.5	27.0	32.0
Secondary	15.4	25.6	18.1	30.6	24.9	41.3
Higher	12.6	17.7	25.2	37.4	29.9	44.8
Wealth quintile						
Lowest	11.9	14.8	12.6	16.3	19.3	24.2
Lower	11.9	17.8	17.4	26.3	23.3	35.2
Middle	15.2	18.8	19.1	23.1	26.0	31.1
Higher	12.8	16.9	22.3	28.1	25.5	32.5
Highest	13.9	17.2	22.4	29.2	31.3	40.3

[1]Married includes married women and those who are living with a man

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

List of indicators for sampling errors, PMA2016/Nasarawa

Variable	Estimation	Base population
Currently using a modern method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using a traditional method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using any contraceptive method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using injectables	Proportion	All women ages 15–49 Married women ages 15–49
Currently using male condoms	Proportion	All women ages 15–49 Married women ages 15–49
Currently using implants	Proportion	All women ages 15–49 Married women ages 15–49
Chose method by self or jointly in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49
Paid fees for family planning services in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49
Informed by provider about other methods	Proportion	All women ages 15–49 Married women ages 15–49
Informed by provider about side effects	Proportion	All women ages 15–49 Married women ages 15–49
Satisfied with provider: Would return and refer friend/relative to provider	Proportion	All women ages 15–49 Married women ages 15–49
Visited by health worker who talked about family planning information in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49

Characteristics of service delivery points

Number and percentage of service delivery points (SDPs) by background characteristics

Background characteristics	Number	Percentage
Total	95	100.0
SDP type		
Hospital	7	7.4
Health center[1]	48	50.5
Pharmacy[2]	39	41.1
Other	1	1.1
Sector		0.0
Public	47	49.5
Private	48	50.5
Offers family planning		
No	9	9.5
Yes	86	90.5

[1]Includes Health Post, Health Clinic, and Maternity Center
[2]Includes Chemist and Patent Medicine Vendors

Charging fees for family planning

Percentage of service delivery points (SDPs) that charge fees for family planning services, by sector and background characteristics, among SDPs offering family planning

Background characteristic	Sector	
	Public	**Private**
Number of SDPs	46	40
Total	26.1	82.5
SDP type		
Hospital	(16.7)	*(100.0)*
Health center[1]	25.6	(83.3)
Pharmacy[2]	na	81.8
Other	*(100.0)*	na

[1]Includes Health Post, Health Clinic, and Maternity Center

[2]Includes Chemist and Patent Medicine Vendors

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Contraceptive choice: Availability of at least 3 or at least 5 modern contraceptive methods

Percentage of public and private service delivery points (SDPs) offering at least 3 or 5 modern contraceptive methods[1], by sector and background characteristics, among all SDPs

SDP Characteristics	3 or more methods		5 or more methods	
	Public	Private	Public	Private
Number	47	48	47	48
Total	74.5	27.1	51.1	2.1
Facility type				
Hospital	(83.3)	*(100.0)*	(83.3)	*(100.0)*
Health Center[2]	72.5	(12.5)	45.0	(0.0)
Pharmacy[3]	na	28.2	na	0.0
Other	*(100.0)*	na	*(100.0)*	na

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Includes Health Post, Health Clinic, and Maternity Center

[3]Includes Chemist and Patent Medicine Vendors

na = not applicable

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Contraceptive choice: Availability of modern contraception, by method

Percentage of service delivery points (SDPs) offering different types of contraceptive methods[1], by sector and method, among all SDPs

	Public SDPs						
	Contraceptive method						
SDP characteristic	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception
Number = 47							
Total	85.1	93.6	34.0	48.9	87.2	2.1	17.0
Facility type							
Hospital	(100.0)	(100.0)	(83.3)	(83.3)	(83.3)	(16.7)	(0.0)
Health center[2]	82.5	92.5	25.0	42.5	87.5	0.0	17.5
Pharmacy[3]	na	na	na	na	na	na	na
Other	*(100.0)*	*(100.0)*	*(100.0)*	*(100.0)*	*(100.0)*	*(0.0)*	*(100.0)*

	Private SDPs						
	Contraceptive method						
SDP characteristic	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception
Number = 48							
Total	58.3	33.3	6.3	6.3	68.8	2.1	27.1
Facility type							
Hospital	*(0.0)*	*(100.0)*	*(100.0)*	*(100.0)*	*(100.0)*	*(100.0)*	*(0.0)*
Health center[2]	(62.5)	(50.0)	(25.0)	(25.0)	(37.5)	(0.0)	(25.0)
Pharmacy[3]	59.0	28.2	0.0	0.0	74.4	0.0	28.2
Other	na	na	na	na	na	na	na

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Includes Health Post, Health Clinic, and Maternity Center

[3]Includes Chemist and Patent Medicine Vendors

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

PMA2020
Performance Monitoring
and Accountability 2020

Contraceptive stock-outs, by method

Percentage of service delivery points (SDPs) stocked out of specific modern contraceptives[1] at time of survey or stocked out in the last 3 months, by sector and background characteristics, among SDPs that offer each method

	Public SDPs					
	Contraceptive method					
Background characteristic	Pill	Injectables	IUD	Implants	Male condom	Emergency contraception
Number of SDPs	40	44	16	23	41	8
Total	20.0	15.9	(18.8)	(30.4)	14.6	(12.5)
SDP type						
Hospital	(33.3)	(50.0)	(20.0)	(40.0)	(20.0)	na
Health center[2]	15.2	10.8	(20.0)	(29.4)	14.3	(14.3)
Pharmacy[3]	na	na	na	na	na	na
Other	*(100.0)*	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*

	Private SDPs					
	Contraceptive method					
Background characteristic	Pill	Injectables	IUD	Implants	Male condom	Emergency contraception
Number of SDPs	28	16	3	3	33	13
Total	21.4	(50.0)	*(33.3)*	*(66.7)*	15.2	(46.2)
SDP type						
Hospital	na	*(0.0)*	*(0.0)*	*(0.0)*	*(0.0)*	na
Health center[2]	*(60.0)*	*(25.0)*	*(50.0)*	*(100.0)*	*(66.7)*	*(100.0)*
Pharmacy[3]	(13.0)	(63.6)	na	na	10.3	(36.4)

[1]Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

[2]Includes Health Post, Health Clinic, and Maternity Center

[3]Includes Chemist and Patent Medicine Vendors

( ) = Estimate generated from less than 25 unweighted observations

( ) = Estimate generated from less than 5 unweighted observations

Other	na	na	na	na	na	na

Number of family planning visits, by method

Total number of visits and new clients for contraceptive services, by method type and background characteristics, among advanced[1] service delivery points (SDPs) that offer each method

	Contraceptive Methods													
	Sterilization		Injectable		IUD		Implants		Male condom		Pills		Emergency Contraception	
	Total women	Total men	Total	New	Total	New	Total	New	Total	New	Total	New	Total	New
Number of SDPs	2	1	49		19		26		45		45		10	
Total	2	0	1,255	271	131	104	473	325	2,733	1,223	504	316	41	10
Background characteristic														
SDP type														
Hospital	2	0	44	2	91	89	37	21	105	19	120	97	0	0
Health center[1]	na	na	1,205	269	40	15	421	289	2,628	1,204	384	219	41	10
Other	na	na	6	0	0	0	15	15	0	0	0	0	0	0
Sector														
Public	0	0	1,217	269	126	102	464	320	2,643	1,220	489	316	35	10
Private	2	na	38	2	5	2	9	5	90	3	15	0	6	0

[1]Includes Health Post, Health Clinic, and Maternity Center